

Algorithm-independent framework for verifying integer constraints*

David Teller[†] Zhong Shao
dtelle@ens-lyon.fr shao@cs.yale.edu
Technical Report YALEU/DCS/TR-1195

July 20, 2000

Abstract

Proof-carrying code (PCC), as pioneered by Necula and Lee, allows a code producer to provide a compiled program to a host, along with a formal proof of safety. The PCC-based systems often rely on solving integer constraints to prove the soundness of the index types and to control resource consumption. Unfortunately, existing approaches often require the inclusion of an oracle-like constraints solver into the trusted computing base (TCB) or at least lock the safety policy with one particular solver. This paper presents a feasibility study for dissociating the constraints solver from the TCB and the safety policy from the actual solver algorithm. To demonstrate this, we produce a simple framework, we show how to adapt the popular solvers such as the Omega test and the Simplex method into this framework and we study some of its properties.

keywords : certified code, certified array bounds checking elimination, certified parallelization, certified Omega test, certified formal calculus, axiomatization of \mathbb{Z}

*This research was sponsored in part by the Defense Advanced Research Projects Agency ISO under the title “Scaling Proof-Carrying Code to Production Compilers and Security Policies,” ARPA Order No. H559, issued under Contract No. F30602-99-1-0519 and in part by NSF Grants CCR-9901011. The views and conclusions contained in this document are those of the authors and should not be interpreted as representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U.S. Government.

[†]This work was done while the first author was visiting the FLINT group in the Department of Computer Science at Yale University.

Contents

1	Introduction	4
2	Background and motivation	5
2.1	Solving integer constraints	5
2.2	Verifying integer constraints	6
2.2.1	Dissociating collection and verification	6
2.2.2	Dissociating the policy from the constraints checker	6
2.2.3	Proving the result	7
3	Solver2FOL	8
3.1	Presentation	8
3.2	Constructs	9
3.2.1	Grammar	9
3.2.2	Semantics	10
3.3	Definitions as axioms	12
3.3.1	First generation	12
3.3.2	Second generation	12
3.4	Basic theorems	14
3.5	General-purpose lemmas	15
3.6	Summary	17
4	The Omega test	17
4.1	The original algorithm	17
4.2	Omega2FOL	18
4.3	An example	22
5	Simplex	22

6	Properties of Solver2FOL	24
6.1	Representation power	24
6.1.1	Introduction	24
6.1.2	Expressing complex functions through brute force	24
6.1.3	Lessons from the brute-force approach	26
6.1.4	Summary	27
6.2	Expressiveness of proofs	27
7	From theory to implementation	27
7.1	Integrating Solver2FOL in a real system	27
7.2	Twelf	28
8	Future work	29
8.1	Implementation	29
8.2	Extensions to non-linear constraints	29
8.3	Exceptions handling	29
9	Related work	30
10	Conclusion	30
A	Proofs using Solver2FOL	32
A.1	Foreword	32
A.2	Syntactical conventions	32
A.3	Basic theorems	32
A.4	The Omega test	52
B	Proofs on Solver2FOL	61
B.1	Expression through Disjunctive Normal Form	61

1 Introduction

In proof-carrying code (PCC) [8, 10], a code producer and a code consumer (host) start by agreeing on a safety policy. This policy is specified as a set of axioms for reasoning about safety. The code producer will then ship a compiled program to the consumer, along with a formal proof of its safety. Of course, the formal proof must be expressed in terms of those axioms. Common examples of such policies include memory soundness, security, CPU time bounds and other resource control. Someday, banking applets will presumably comply with money-transfer soundness policies.

PCC relies on the same formal methods as does program verification; but it has the significant advantage that safety properties are much easier to prove than program correctness. The producer's formal proof will not, in general, prove that the code produces a correct or meaningful result; but it guarantees that execution of the code can do no harm. Thus, it cannot replace other methods of program assurance. On the other hand, the proofs can be mechanically checked by the host; the producer need not be trusted at all, since a valid proof is incontrovertible evidence of safety.

Using PCC allows to remove many run-time checks without sacrificing safety. For example, the Touchstone compiler [10] can prove the memory safety of the compiled programs. In other words, Touchstone-compiled programs can be trusted to run on devices without memory protection. Recently, Xi [19, 17] introduced a dependent type system in which the costly process of array bounds checking can be removed—a method which has been adapted to do so in a provably safe way.

CPU time bounding [9, 5], memory soundness [8, 7, 1], and array bounds checking [19], all require some kind of integer constraints handling. So does automatic parallelization [14]. And so do presumably many unmentioned processes. To be more specific, they require solving a set of integer equality or inequality equations to prove the soundness of the index types and to control resource consumption. Unfortunately, existing approaches either include the constraints solver into the trusted computing base (e.g., Xi's DML [17] and DTAL [18]), or lock the safety policy with one particular solver logic (e.g., Necula's Simplex solver logic [10, 9]).

The goal of this work is to study the feasibility of producing a generic framework where the constraints solver does not have to be in the TCB and the safety policy is independent of the actual solver algorithm. This is important because it would allow the code producer to choose the constraints solver most appropriate to a particular application. The producer will have much more flexibility in the manner in which the mobile code is proved safe. Furthermore, by removing the solver algorithm or the specific solver logic inference rules from the TCB, we get a higher-assurance system: any assumption about a particular solver algorithm must be proved.

2 Background and motivation

2.1 Solving integer constraints

Trying to solve all integer constraints is a somewhat ambitious task. As a matter of fact, this would mean solving Hilbert's tenth problem "Determination of the solvability of a Diophantine equation. Given a diophantine equation with any number of unknown quantities and with rational integral numerical coefficients: To devise a process according to which it can be determined by a finite number of operations whether the equation is solvable in rational integers." A problem which has been showed undecidable.

Hence, no solver handles just any integer constraints. Most of them only handle linear integer constraints. Some of them contain extensions to simple families of polynomials. We will follow the wise precedents and only attack linear constraints, for now.

Three families of solvers are currently used :

- Fourier-Motzkin's variable elimination [6]
- SUP-INF [3]
- The Simplex algorithm [11]

The most commonly used algorithm is the Omega test [13], a variant on Fourier-Motzkin's variable elimination [6]. Supposedly, it is the fastest and most complete algorithm currently in use. For now, suffice to say that it works by eliminating trivial equalities and inequalities, reducing non-trivial equalities into trivial equalities by applying a variant of the mod operator, reducing non-trivial inequalities into trivial equalities by projection in some n-dimensional space and checking exhaustively for solutions when the formal methods do not work. Since this is the most commonly used algorithm and this it is definitely non-trivial, we used it as a basis for our research.

The Simplex algorithm, noticeably used by Necula in [Necula 1998], is a much simpler algorithms which proceeds by some linear algebra transformations on the matrix representation of the constraints. Since this algorithm is fairly simple, completely different from the Omega test, and since Necula has already made some steps to make it produce proofs, we used it as a confirmation of the genericity of this work.

2.2 Verifying integer constraints

2.2.1 Dissociating collection and verification

The first approach one can think of for the verification of integer constraints-related policies is the definition of a source-level semantic/logic. In other words, a set of inference rules which would be applied directly to the source code, would check all integer-related operations and decide, for example, of the satisfiability of problematic cases. This verification could take, for example, the form of a type-checking algorithm.

A variant of this approach would imply the translation into a simpler language, say some form of λ -calculus, which would allow the use of a smaller set of rules.

The shortcoming of this method is in its definition : interleaved collection and resolution of integer constraints - both of which are undecidable problems - make this semantic bigger, more complex, harder to maintain. And, of course, if the security policy is to change, the whole semantic will be due for change.

Dissociating both aspects is not going to solve anything by itself, of course. But it allows to work on both aspects separately, and to find separate solutions. For example, using Xi's dependent type system [17] or an annotation system for the collection of integer constraints makes it possible to overcome the undecidability of constraints collection.

2.2.2 Dissociating the policy from the constraints checker

The logical successor of the former approach is a two-level architecture : one constraints collector and one policy checker/prover. However, this is still no solution.

Some program require different security policies. For example, two different optimizations such as array bounds checking elimination and automatic parallelization may not be proved valid by the same set of criteria. A program taking advantage of both optimizations will hence be collected for constraints and then policy-checked/policy-proved twice. However, both policies rely on integer constraints verification. In other words, it might be that only one step of both policy solver/checker is different.

In other words, the size of the proof and the time required to produce and then check it can be reduced by just dissociating the policy checker and the constraints solver. For example, the Omega Test [13] is an "as generic as it gets" integer constraints solver.

2.2.3 Proving the result

The next step is that of solving the constraints, which can be handled by classical algorithms such as the Omega Test or the Simplex for given families of integer constraints.

This is, however, not the last step. For the receiving host still has to be convinced of the (un)satisfiability of the set of constraints.

The satisfiability is rather trivial to prove, since it only requires a numerical certificate. In other words, a numerical example. However, the unsatisfiability is a much harder problem. In practice, when the enforcement of the policy requires the unsatisfiability of a set of constraints, two distinct methods are used to check that the security policy is enforced.

2.2.3.1 Solver algorithm in TCB The first solution is to include the decision algorithm in the trusted computing base. In other words, the run-time proof checking system contains the Omega test, for example, and is submitted a set of linear equations and inequations, which it solves, just like the compiler did. The method seems to have been implicitly used by Xi [17].

This method is not without advantages :

- It is simple to implement.
- The included Omega test might be very low level and fast.
- Carried proofs can be very short.

However, it has serious drawbacks :

- Since the whole set of equations and inequations is solved again, it requires lots of possibly useless calculations.
- The algorithm used cannot be changed, which prevents, for example, from adding the capability of solving some simple kind of polynomial constraints, to the solver.
- The TCB must be expanded with a possibly complex algorithm, thus increasing the risks of bugs, leading to unsoundness.
- If the run-time system has to be ported to another platform, the included test makes it bigger, hence probably harder to port.
- Since no integer constraints solving algorithm is complete, some kinds of constraints cannot be checked at compile-time. In particular, if the type system requires these constraints, as is the case with dependent/indexed type systems [17], the compiler will flatly refuse perfectly sound programs.

2.2.3.2 Solver-logic in TCB Another simple idea, introduced by Necula [10], is to define a set of deduction/inference rules encompassing all the steps in the algorithm and proving the result using these rules. In fact, this is the same thing as introducing a new logic adapted to the solver.

Once again, this has advantages:

- The algorithm does not need to be included in the TCB as such.
- Some useless calculations can be removed from the proof, hence resulting in faster checks.

However, similar drawbacks can be found:

- The TCB is still expanded, this time with a specific logical system, which is almost as hard to trust as the algorithm itself.
- The system is still locked with a particular algorithm and minor variations on this algorithm. This approach still prevents some optimizations and still refuses some sound programs.

3 Solver2FOL

3.1 Presentation

One of the next logical steps of modularization in the verification of integer constraints is that of rendering the algorithm independent from the proof. That of getting rid of the limits of the approaches previously exposed.

Solver2FOL is a small contribution to this task: a study of the feasibility of this dissociation. Although this work is no way related to the semantic model for PCC introduced by Appel and Felty [1], it can be seen as a complementary module: where Appel and Felty introduce a semantic model to describe types and machine instructions, in order to achieve language independence in PCC, we introduce a model to describe constraints solvers, in order to achieve algorithm independence in PCC.

Solver2FOL is a set of tools designed to allow the translation of integer constraints solving algorithms into algorithms building First Order Logic (FOL) proofs. It is composed of :

- A syntax to express constraints.
- A minimal semantics on the constraints.
- A minimal list of trusted operations.

- A collection of formal definitions to enrich the constraints using usual comparisons and operations.
- A collection of basic theorems, all of them formally proved, which prove the use of these enrichments valid.
- A collection of general-purpose lemmas, formally proved using these theorems.

In addition, we provide as an example Omega2FOL and Simplex2FOL, formally proved “Solver2FOL-izations” of the Omega test and of the Simplex.

3.1.0.3 FOL vs. set of rules First Order Logic was chosen instead of a custom set of rules since First Order Logic allows to express the kind of rules such a work would need as a set of axioms. Using a custom set of rules would provide shorter proofs but would not allow us to use the full power of First Order Logic. Unless, of course, we want to reexpress FOL into this set of rules.

And of course, using a custom set of rules is the first step toward defining and locking the algorithm by the mere description of the proof system - which is exactly what we want to avoid. FOL allows us to ignore this pitfall.

3.1.0.4 FOL vs. HOL First Order Logic was chosen because FOL rules and axioms tend to be simpler and more readable than true Higher Order Logics. However, we will show that this approach has several drawbacks.

3.1.0.5 Formalism Solver2FOL redefines most operations on \mathbb{Z} using formal notations. This redefinition allows us to produce *purely formal proofs* of lemmas and theorems. In time, these purely formal proofs may be shipped and mechanically verified by a theorem checker.

3.2 Constructs

3.2.1 Grammar

Figure 1 gives the grammar for Solver2FOL. *Tests* are to be taken as *First Order Logic* terms.

This grammar is chosen as to make the basic set of axioms be as minimal as possible. In particular, the grammar does not contain inequalities or division—all these can be built on top of this skeleton.

This grammar is also designed to be fairly generic, so it can express the expressions as seen by the solver algorithm. For example, the Omega test mostly works with expressions of form

$Test ::= Term \approx Term$ (term comparison)

$Term ::= Var \in \mathcal{V}$ (unknown variable)
| $Int \in \mathbb{Z}$ (integer constant)
| $Term \oplus Term$ (term addition)
| $Term \otimes Term$ (term multiplication)

Where \mathcal{V} is a countable infinite set.

Figure 1: Grammar for Solver2FOL

“ $Term \preceq 0$ ” along sometimes with “ $a \otimes z \preceq \alpha$ ” and “ $\beta \preceq b \otimes z$ ”, whereas the Simplex only uses “ $Term \succeq 0$ ” and “ $Term \approx Term$ ”.

These design decisions allow the translated Solver2FOL algorithm to use the exact same set of conditions as the original algorithm as well as to reflect directly its inner workings. The other reasons for our choice are to serve as a place-holder for future extensions on polynomials and to pave the way to render the translation from an algorithm into Solver2FOL semi-automatic.

3.2.2 Semantics

3.2.2.1 Constants

Comparisons on constants Classical comparisons on \mathbb{Z} such as \leq , $<$, $=$, $>$, \geq and $|$ are supposed to be already implemented and trusted. This represents the fact that all those operations are already part of the CPU, hence already trusted in order to run the proof-checking system.

Operations on constants Classical operations on \mathbb{Z} such as addition, multiplication, floor division ($\lfloor \cdot \rfloor$), ceil division ($\lceil \cdot \rceil$), modulo and absolute value are supposed to be already implemented and “mostly trusted”. This represents the fact that all those operations are already part of the CPU, hence already trusted in order to run the proof-checking system - but that exceptions can occur.

“Mostly trusted” means that operations are trusted as long as no exceptions are thrown (division by zero, overflow, underflow). The semantics of exceptions in Solver2FOL is that if any exception should be raised, then the whole proof is considered false. Although this is not a fully satisfactory arrangement, it is not incoherent : if a proof contains such an exception, it

probably means that the prover forgot to catch the same exception. However, this might not be true in case of cross-platform proof-building.

3.2.2.2 Variables

Operations on variables Although most of the times integer constants will not be effectively substituted to variables, the semantics of Solver2FOL is designed for “constant place-holders”-variables.

In other words, variables stand for unknown constants.

3.2.2.3 Terms

Definition Terms are members of set \mathcal{T} defined by the grammar of Solver2FOL. Terms can be considered as an extension of integers with integer variables. For the sake of simplicity, we will consider \mathcal{T} as a superset of \mathbb{Z} .

Comparisons on terms Classical comparisons on \mathbb{Z} such as \leq , $<$, $=$, $>$, \geq and $|$ can be extended to terms. However, the only needed comparison is the equality law \approx , defined by

$$\left\{ \begin{array}{l} \approx \text{ is an equivalence law on} \\ \mathcal{T} \\ \approx \text{ is compatible with } = \text{ on} \\ \mathbb{Z} \end{array} \right.$$

Operations on terms Classical operations on \mathbb{Z} such as addition, multiplication, division ($[\cdot]$) can be extended to terms. However, the only needed operations are addition \oplus and multiplication \otimes , defined by

$$\left\{ \begin{array}{l} \oplus \text{ is an internal composition law on} \\ \mathcal{T} \\ \oplus \text{ is commutative, associative, dis-} \\ \text{tributive on } \otimes \\ 0 \text{ is neutral for } \oplus \\ \oplus|_{\mathbb{Z}} \equiv_{def} + \end{array} \right.$$

$$\left\{ \begin{array}{l} \otimes \text{ is an internal composition law on } \\ \mathcal{T} \\ \otimes \text{ is commutative, associative} \\ 0 \text{ is absorbant for } \otimes \\ 1 \text{ is neutral for } \otimes \\ \otimes|_{\mathbb{Z}} \equiv_{def} \cdot \end{array} \right.$$

3.3 Definitions as axioms

3.3.1 First generation

3.3.1.1 Syntactical conventions

- Operators on \mathcal{T} have the same precedence as their counterparts in \mathbb{Z} .
- In case of ambiguity, all operations are supposed left-parenthesized.
- $X \ominus Y$ is a syntactical shortcut for $X \oplus ((-1) \otimes Y)$
- lowercase letters stand for integers constants
- uppercase letters stand for expressions

3.3.1.2 Presentation The basic axioms are presented in Figure 2. They are translations of the semantics of the basic constructs of Solver2FOL.

3.3.2 Second generation

3.3.2.1 Definition Operators $\cdot \setminus \cdot$ and $\cdot \% \cdot$ are the respective counterparts of $\lfloor \cdot \rfloor$ and $\cdot \bmod \cdot$ on \mathcal{T} . Binary relations $\parallel, \preceq, \prec, \succ, \succeq$ are the respective counterparts of $|, \leq, <, >, \geq$. They are defined by :

Equality		
$\forall X$	$X \approx X$	(equ ref)
$\forall X, Y$	$X \approx Y \Rightarrow Y \approx X$	(equ sym)
$\forall X, Y, Z$	$X \approx Y \wedge Y \approx Z \Rightarrow X \approx Z$	(equ trans)
$\forall a, b$	$a \approx b \Rightarrow a = b$	(equ on \mathbb{Z})
$\forall X$	$\exists a, X \approx a$	(equ int)
Addition		
$\forall X$	$X \oplus 0 \approx X$	(add 0)
$\forall X, Y$	$X \oplus Y \approx Y \oplus X$	(add comm)
$\forall X, Y, Z$	$X \oplus (Y \oplus Z) \approx (X \oplus Y) \oplus Z$	(add assoc)
$\forall X, Y, Z$	$X \approx Y \Rightarrow X \oplus Z \approx Y \oplus Z$	(add eq)
$\forall a, b$	$a \oplus b \approx a + b$	(add on \mathbb{Z})
Multiplication		
$\forall X$	$0 \otimes X \approx 0$	(mult 0)
$\forall X$	$1 \otimes X \approx X$	(mult 1)
$\forall X, Y$	$X \otimes Y \approx Y \otimes X$	(mult comm)
$\forall X, Y, Z$	$X \otimes (Y \otimes Z) \approx (X \otimes Y) \otimes Z$	(mult assoc)
$\forall X, Y, Z$	$(X \oplus Y) \otimes Z \approx (X \otimes Z) \oplus (Y \otimes Z)$	(add/mult dist)
$\forall X, Y, Z$	$X \approx Y \Rightarrow X \otimes Z \approx Y \otimes Z$	(mult eq)
$\forall a, b$	$a \otimes b \approx a \cdot b$	(mult on \mathbb{Z})

Figure 2: Basic axioms for Solver2FOL

$$\left\{ \begin{array}{l} X \setminus a \approx Y \text{ iff for some } b \text{ in } [0, a - 1], X \approx \\ a \otimes Y \oplus b \\ X \% a \approx b \text{ iff } b \text{ is in } [0, a - 1] \text{ and for some } Y, \\ X \approx a \otimes Y \oplus b \\ a \parallel X \text{ iff } X \% a \approx 0 \\ X \preceq Y \text{ iff for some non-negative } b, Y \approx X \oplus b \\ X \prec Y \text{ iff for some strictly positive } b, Y \approx \\ X \oplus b \\ X \succeq Y \text{ iff for some non-negative } b, X \approx Y \oplus b \\ X \succ Y \text{ iff for some strictly positive } b, X \approx \\ Y \oplus b \end{array} \right.$$

3.3.2.2 Axiomatization The axiomatization of these operators and relations is immediate. It is presented on figure 3.

Definition of operators

$$\begin{aligned} \forall X, Y, a \quad X \setminus a \approx Y &\equiv_{def} \exists b, \\ &0 \leq b \wedge b < a \wedge X \approx a \otimes Y \oplus b & \text{(dodiv)} \\ \forall X, a, b \quad X \% a \approx b &\equiv_{def} \exists Y, \\ &0 \leq b \wedge b < a \wedge X \approx a \otimes Y \oplus b & \text{(mod)} \end{aligned}$$

Definition of relations

$$\begin{aligned} \forall X, a \quad a \parallel X &\equiv_{def} X \% a \approx 0 & \text{(isdiv)} \\ \forall X, Y \quad X \preceq Y &\equiv_{def} \exists b, 0 \leq b \wedge Y \approx X \oplus b & \text{(leq)} \\ \forall X, Y \quad X \prec Y &\equiv_{def} \exists b, 1 \leq b \wedge Y \approx X \oplus b & \text{(lt)} \\ \forall X, Y \quad X \succeq Y &\equiv_{def} \exists b, 0 \leq b \wedge X \approx Y \oplus b & \text{(geq)} \\ \forall X, Y \quad X \succ Y &\equiv_{def} \exists b, 1 \leq b \wedge X \approx Y \oplus b & \text{(gt)} \end{aligned}$$

Figure 3: The second wave of definitions

3.4 Basic theorems

The following set of theorems will prove that the newly introduced operators are what they seem - namely that they can be used as their integer counterparts. Purely formal proofs are given in appendix.

Basic theorem 1 (*dodiv on \mathbb{Z}*) *The definition of \setminus is compatible with $\lfloor \cdot \rfloor$ on \mathbb{Z} . In other words, for any integers a, b , $a \setminus b \approx \lfloor \frac{a}{b} \rfloor$.*

Basic theorem 2 (*mod on \mathbb{Z}*) *The definition of $\%$ is compatible with $\cdot \bmod \cdot$ on \mathbb{Z} . In other words, for any integers a, b , $a \% b \approx a \bmod b$.*

Basic theorem 3 (*isdiv on \mathbb{Z}*) *The definition of \parallel is compatible with $|$ on \mathbb{Z} . In other words, for any integers a, b , $a \parallel b \iff a|b$*

Basic theorem 4 (*leq on \mathbb{Z}*) *The definition of \succeq is compatible with \geq on \mathbb{Z} . In other words, for any integers a, b , $a \succeq b \iff a \geq b$*

Basic theorem 5 *\succeq defines a partial order on \mathcal{T} . In other words,*

$$\left\{ \begin{array}{ll} \forall X & X \succeq X & \text{(geq ref)} \\ \forall X, Y & X \succeq Y \wedge Y \succeq X \Rightarrow X \approx Y & \text{(geq antis)} \\ \forall X, Y, Z & X \succeq Y \wedge Y \succeq Z \Rightarrow X \succeq Z & \text{(geq trans)} \end{array} \right.$$

Basic theorem 6 (*gt on \mathbb{Z}*) The definition of \succ is compatible with $>$ on \mathbb{Z} . In other words, for any integers a, b , $a \succ b \iff a > b$

Basic theorem 7 (*lt on \mathbb{Z}*) The definition of \prec is compatible with $<$ on \mathbb{Z} . In other words, for any integers a, b , $a \prec b \iff a < b$

Basic theorem 8 (*leq on \mathbb{Z}*) The definition of \preceq is compatible with \leq on \mathbb{Z} . In other words, for any integers a, b , $a \preceq b \iff a \leq b$

Basic theorem 9 \preceq defines a partial order on \mathcal{T} . In other words,

$$\begin{cases} \forall X & X \preceq X & (\text{leq ref}) \\ \forall X, Y & X \preceq Y \wedge Y \preceq X \Rightarrow X \approx Y & (\text{leq antis}) \\ \forall X, Y, Z & X \preceq Y \wedge Y \preceq Z \Rightarrow X \preceq Z & (\text{leq trans}) \end{cases}$$

Basic theorem 10 (*leq equiv geq*) \preceq and \succeq are each other's symmetric. In other words, $\vdash X \preceq Y \iff Y \succeq X$.

Basic theorem 11 (*lt equiv gt*) \prec and \succ are each other's symmetric. In other words, $\vdash X \prec Y \iff Y \succ X$.

Basic theorem 12 (*lt as leq*) $\preceq = \prec \cup \approx$. In other words, $\vdash X \preceq Y \iff (X \prec Y \vee X \approx Y)$.

Basic theorem 13 (*gt as geq*) $\succeq = \succ \cup \approx$. In other words, $\vdash X \succeq Y \iff (X \succ Y \vee X \approx Y)$.

Basic theorem 14 (*eq as ineq*) $\preceq \cap \succeq = \approx$. In other words, $X \preceq Y \wedge X \succeq Y \vdash X \approx Y$.

Basic theorem 15 (*absurd ineq*) $\prec \cap \succ = \emptyset$. In other words, $X \prec Y \wedge X \succ Y \vdash \perp$.

3.5 General-purpose lemmas

Formal proofs are given in appendix.

Main lemma 1 (*opp add*) Opposite addition is provable using Solver2FOL. In other words, $\vdash Y \ominus Y \approx 0$

Main lemma 2 (*add eq2*) A second formulation for addition on each side of an equality can be proved using Solver2FOL. In other words, $X \approx Y \vdash Z \oplus X \approx Z \oplus Y$

Main lemma 3 (*mult eq2*) A second formulation for addition on each side of an equality can be proved using Solver2FOL. In other words, $X \approx Y \vdash Z \otimes X \approx Z \otimes Y$

Main lemma 4 (*add eq3*) General addition on both side of the equality sign can be proved using Solver2FOL. In other words, $X \approx Y \wedge Z \approx T \vdash X \oplus Z \approx Y \oplus T$

Main lemma 5 (*subtrac equ*) Substraction in equality (*sub equ*) is provable using Solver2FOL. In other words,

$$\begin{cases} X \oplus Y \approx Z \vdash X \approx Z \ominus Y \\ Z \approx X \oplus Y \vdash Z \ominus Y \approx X \end{cases}$$

Main lemma 6 (*sub ineq*) Substraction in an inequality is provable with Solver2FOL. In other words,

$$X \oplus Y \preceq Z \vdash X \preceq Z \ominus Y$$

Main lemma 7 (*mult sum*) The propagation of multiplication in a sum is provable with Solver2FOL. In other words,

$$\vdash p \otimes \Sigma_{i \in I} X_i \approx \Sigma_{i \in I} p \otimes X_i$$

Main lemma 8 (*var isol*) The isolation against a variable in a sum is provable with Solver2FOL. In other words, for any j in I ,

$$\vdash \Sigma_{i \in I} X_i \approx (\Sigma_{i \in I \setminus \{j\}} X_i) \oplus X_j$$

Main lemma 9 (*var resol*) The resolution against a variable of an equality between a sum and a constant is provable in Solver2FOL. In other words, for any k in I ,

$$\Sigma_{i \in I} X_i \approx c \vdash X_k \approx c \oplus \Sigma_{i \in I \setminus \{k\}} \ominus X_i$$

Main lemma 10 (*0 add2*) $\vdash X \approx X \oplus 0$

Main lemma 11 (*div sum*) The propagation of $\cdot \setminus p$ in a sum where all elements are divisible by p is provable using Solver2FOL. In other words, if $p|a_1, p|a_2, \dots, p|a_n$ then $\vdash (\Sigma_{i \in [1, n]} a_i \otimes X_i) \setminus p \approx \Sigma_{i \in [1, n]} \frac{a_i}{p} \otimes X_i$

Main lemma 12 (*mult ineq*) Multiplication in inequality is provable using Solver2FOL. In other words,

$$X \preceq Y \wedge 0 \preceq Z \vdash Z \otimes X \preceq Z \otimes Y$$

3.6 Summary

Once this set of theorems and lemmas is proved, we have a full syntax and its associated semantics to handle formal expressions. Note that as opposed to Xi's approach, for example, quantifiers are not handled by Solver2FOL - they don't need to, since they are already handled by First Order Logic.

Also note that the trusted computing base is very small: it consists of classical integer operations and very few (and very basic) properties of \otimes and \oplus . Thus, complex proofs can be traced back to a base of axioms which can easily be checked and trusted.

4 The Omega test

4.1 The original algorithm

The input of the Omega test is a set of linear equalities and linear inequalities involving only integer values. All the conditions are supposed to have form $\sum_{i=1}^{i=n} a_i \cdot x_i \leq c$ or $\sum_{i=1}^{i=n} a_i \cdot x_i = c$.

4.1.0.3 Removing equality constraints

- Equalities of the form $\sum_{i=1}^{i=n} a_i \cdot x_i = c$ are divided by the greatest common divisor of a_1, \dots, a_n . If this leads to a formula with non-integer values, the test has failed and the system is unsatisfiable.
- If two equalities are visibly incompatible, the set is unsatisfiable.
- If one of the equalities contains a variable with coefficient 1 or -1, remove this equality and eliminate the variable.
- Otherwise, choose one equality $\sum_{i=1}^{i=n} a_i \cdot x_i = c$ and replace it with $-|a_k \cdot x_k| \cdot \sigma + \sum_{i=1}^{i=n, i \neq k} (\lfloor a_i/m + 1/2 \rfloor + a_i \mu m) \cdot x_i$ where $a \mu b \equiv_{def} a - b \cdot \lfloor a/b + 1/2 \rfloor$ and $m \cdot \sigma \equiv_{def} \sum_{i=1}^{i=n} (a_i \mu m) \cdot x_i = c$
- Proceed until there are no more equality constraints.

4.1.0.4 Removing inequality constraints

- Inequalities of the form $\sum_{i=1}^{i=n} a_i \cdot x_i \leq c$ are divided by the greatest common divisor of a_1, \dots, a_n , rounding down c/g .
- If inequalities can be merged into equalities, merge them.
- If two inequalities are redundant, remove the weakest one.

- If inequalities are visibly incompatible, the set is unsatisfiable.
- If there are no more equalities and if there is a variable without upper bound or without lower bound, remove all the inequations involving this variable.
- Compute the *real shadow* of the set of inequalities along z by merging two constraints such as $a \cdot z \leq \alpha$ and $\beta \leq b \cdot z$ into constraint $a \cdot \beta \leq b \cdot \alpha$, where a and b are positive integers.
- Compute the *integer shadow* of the set of inequalities along z by merging the two same constraints into $b \cdot \alpha - a \cdot \beta \geq (a - 1) \cdot (b - 1)$.
- If both shadows are identical, the system is satisfiable iff there are integer solutions to the (common) shadow
- otherwise
 - if there are no integer solutions to the real shadow, the set is unsatisfiable
 - if there are integer solutions to the integer shadow, the set is satisfiable
 - otherwise
 - * determine the largest coefficient a of z in any upper bound $\alpha \geq az$ on z and for each lower bound $bz \geq \beta$ on z , for each i in $[0, (ab - a - b)/a]$ do solve the problem combined with $bz = \beta + i$
 - Proceed until there is at most one inequality. If this happens the system is satisfiable. Otherwise, it is not.

4.2 Omega2FOL

Let us consider a reduced version Omega' of the known Omega test. This Omega' is a subset of the Omega test where two optimizations have been remove for the sake of simplicity. These optimizations are not required to run the Omega test :

Coefficients reduction We have not tried to prove this step yet. However, our work so far seems to indicate that this step will require a more complete work on the Omega test, including completeness studies.

Fusion of opposite inequalities This step is trivial but meaningless without coefficients reduction.

Construct 1 *There exists an algorithm Omega2FOL based on Solver2FOL which :*

- *can solve the same set of constraints as Omega'.*
- *instead of Yes/No, returns **proofs of unsatisfiability** whenever Omega' decides that the set is unsatisfiable.*

4.2.0.5 Steps All these steps have been formally proved using Solver2FOL. Proofs are presented in annex.

Algorithm step 1 “Equality normalization” is provable using Solver2FOL. In other words, if $p|a_1, p|a_2, \dots, p|a_n$ then $\sum_{i \in [1, n]} a_i \otimes X_i \approx c \vdash \sum_{i \in [1, n]} \frac{a_i}{p} \otimes X_i \approx \lfloor \frac{c}{p} \rfloor$

Algorithm step 2 “Unsatisfiable equality” is provable using Solver2FOL. In other words,

$$p|a_1 \wedge p|a_2 \wedge \dots \wedge p|a_n \wedge \sum_{i=1}^n a_i \otimes X_i \approx c \vdash p|c$$

Algorithm step 3 “Inequality tightening” is provable using Solver2FOL. In other words, if $p|a_1, p|a_2, \dots, p|a_n$, then

$$\sum_{i \in I} a_i \otimes X_i \preceq c \vdash \sum_{i \in I} \frac{a_i}{p} \otimes X_i \preceq \lfloor \frac{c}{p} \rfloor$$

Algorithm step 4 “Real shadow” can be proved using Solver2FOL. In other words, if $1 \leq a$, $1 \leq b$ then $a \otimes Z \preceq A \wedge B \preceq b \otimes Z \vdash a \otimes B \preceq b \otimes A$.

Algorithm step 5 “Exhaustive check” can be proved using Solver2FOL. In other words, if $1 \leq a \wedge 1 \leq b$, $(a - 1) \cdot (b - 1) \preceq b \otimes A \ominus a \otimes B \wedge a \otimes \xi \leq A \wedge B \leq b \otimes \xi \vdash \exists i, 0 \leq i \wedge i \cdot a \leq (a \cdot b - a - b) \wedge b \otimes \xi \approx B \oplus i$

Corollaries :

- Omega’ was proved.
- We can produce Omega2FOL.

4.2.0.6 Informal listing of Omega2FOL For the sake of readability, this listing is an informal presentation of the decision procedure. For example, it does not detail the instantiation of each lemma, nor does it explicitly show that it tries to simplify the handled equations at each step, by turning multiplication by 0 into 0, by removing addition of 0, etc...

Also note that this algorithm returns a formal proof where Omega’ would have returned *No* and *Maybe* where Omega’ would have returned *Yes*. This seemed more appropriate in absence of any completeness study on Omega’.

Removing equality constraints

If there is at least one unnormalized equality left \mathcal{E}

Normalize

Call Omega2FOL with a system containing the normalized equality \mathcal{F} instead of the unnormalized one.

If the test returns *Maybe*, return *Maybe*

Otherwise, if the test returns proof P

Return the composition of proof $\vdash \mathcal{E} \Rightarrow \mathcal{F}$ (built by lemma “Equality normalization”) and P

If we can find an unsatisfiable equality \mathcal{E}

Return lemma “Unsatisfiable equality” applied to \mathcal{E}

If we can find two contradictory equalities A and B

Return lemma “eq trans” applied to A and B

If there is one equality \mathcal{E} which involves a variable ξ with coefficient 1

Remove one variable

Let $(\mathcal{F}_i)_i$ be the set of equations and inequations. Call Omega2FOL with a system containing $(\mathcal{F}'_i)_i$, the set of all equalities/inequalities in which the formal value associated to ξ has been substituted to ξ itself.

If the test returns *Maybe*, return *Maybe*.

Otherwise, if the test returns proof P

Return the composition of the proof of the formal value of ξ (built by lemma “variable resolution”), the proofs $\vdash \mathcal{F}_i \Rightarrow \mathcal{F}'_i$ (built by lemma “variable isolation in a sum” and lemma “add eq2”) and P.

If there is one equality \mathcal{E} which involves a variable with coefficient -1

Remove one variable

Call Omega2FOL with a system containing the opposed equality \mathcal{F} in which the -1 has been fully propagated.

If the test returns *Maybe*, return *Maybe*

Otherwise, if the test returns proof P

Return the composition of proof $\vdash \mathcal{E} \Rightarrow \mathcal{F}$ (built by lemma “eq mult” and lemma “sum mult”) and P

Otherwise, if there is still at least one equality, let $X \approx c$ be one of the equalities left

Call Omega2FOL with a system containing $X \preceq c$ and $c \succeq X$ instead of $X \approx c$

If the test returns *Maybe*, return *Maybe*

Otherwise, if the test returns proof P

Return the composition of proof $\vdash X \approx c \Rightarrow X \preceq c \wedge X \succeq c$ (built by lemma “eq as ineq”) and P

Removing inequality constraints

If we can find two redundant constraints

Call Omega2FOL with a system containing all the constraints minus the weakest of the two redundant constraints. Return the result.

If we can find two contradictory inequalities A and B

Return lemma “leq trans” applied to A and B

If there are no more equalities and if we can find a variable without upper bound or without lower bound.

Call Omega2FOL with a system containing only comparisons which do not involve this variable. Return the result.

If there is a variable ζ , along with one lower bound condition $B \preceq b \otimes \zeta$ and one upper bound condition $a \otimes \zeta \preceq A$ such as $a = 1$ and $1 \leq b$ or $b = 1$ and $1 \leq a$.

Both shadows are identical

Call Omega2FOL with a system containing $a \otimes B \preceq b \otimes A$ instead of both conditions

If the test returns *Maybe*, return *Maybe*

Otherwise, if the test returns proof P

Return the composition of proof $\vdash B \preceq b \otimes \zeta \wedge a \otimes \zeta \preceq A \Rightarrow a \otimes B \preceq b \otimes A$ (built by lemma “Real shadow”) and P

If there is a variable ζ , along with one lower bound condition $B \preceq b \otimes \zeta$ and one upper bound condition $a \otimes \zeta \preceq A$ ($a, b \in \mathbb{N}^*$). Ideally, choose the largest a and b available.

Shadows are not identical

Call Omega2FOL with a system containing $a \otimes B \preceq b \otimes A$ instead of both conditions

If the test returns *Maybe*

Call Omega2FOL with a system containing $(a - 1) \cdot (b - 1) \preceq b \otimes A \ominus a \otimes B$ instead of both conditions

If the test returns *Maybe*, return *Maybe*

Otherwise, if the test returns proof P

Try all possibilities

for each i in $[0, \frac{ab-a-b}{a}]$ do

Call Omega2FOL with a system containing $b \otimes \zeta \approx B \oplus i$ instead of $B \preceq b \otimes \zeta$

If the test returns *Maybe*, return *Maybe*
 Otherwise, let P_i be the proof returned by the test
 Return the composition of $(P_i)_i$ and lemma “Exhaustive check” composed
 itself with P
 return the proof

Otherwise, if the test returns proof P
 Return the composition of P and lemma “Real shadow”

If there is only one condition left, and if it is an inequality involving a variable, return *Maybe*

4.3 An example

Consider the system

$$\left\{ \begin{array}{l} X \ominus Y \approx 0 \\ X \oplus Y \approx 0 \\ X \preceq -1 \\ \mathcal{S} \equiv_{def} X \ominus Y \approx 0 \wedge X \oplus Y \approx 0 \wedge X \preceq -1 \end{array} \right.$$

If we ignore the omnipresent “eq trans” axiom (transitivity of equality), the resolution will look like figure 4

5 Simplex

The Simplex algorithm is a well known integer constraints solving algorithm. Noticeably, it has been used by Necula and Lee [10] in order to produce proofs in the Touchstone compiler.

Using Necula’s work, the Simplex can be seen as a decision procedure building proofs using 9 simple logical rules : the Simplex-logic rules, presented on figure 5. Of these rules, *geqgeq*, *gtgeq*, *leqgeq*, *ltgeq*, *eqgeq*, *geq0* are already axioms or lemma of Solver2FOL and the three other ones can be very easily translated to Solver2FOL.

Construct 2 *There exists an algorithm Simplex2FOL based on Solver2FOL which :*

- *can solve the same set of constraints as the Simplex.*
- *instead of Yes/No, returns **proofs of unsatisfiability** whenever the Simplex decides that the set is unsatisfiable.*

$X \ominus Y \approx 0$ is an equality involving variable X with coefficient 1
Resolution of the equality against X
 $\mathcal{S} \vdash X \approx Y$ (proved by lemma “variable resolution”)
Introduction of the value of X into $X \oplus Y \approx 0$
 $\mathcal{S} \vdash Y \oplus X \approx 0$ (proved by lemma “variable isolation”)
 $\mathcal{S} \vdash Y \oplus Y \approx 0$ (proved by axiom “add eq”)
Introduction of the value of X into $X \preceq -1$
 $\mathcal{S} \vdash Y \preceq -1$ (proved by “eq as ineq”)
Factorization of $Y \oplus Y \approx 0$
 $\mathcal{S} \vdash 1 \otimes Y \oplus Y \approx 0$ (proved by “add eq” and “mult 1”)
 $\mathcal{S} \vdash 1 \otimes Y \oplus 1 \otimes Y \approx 0$ (proved by “add eq” and “mult 1”)
 $\mathcal{S} \vdash 2 \otimes Y \approx 0$ (proved by “mult/add dist”)
Equality normalization of $2 \otimes Y \approx 0$
 $\mathcal{S} \vdash Y \approx 0$ (proved by “equality normalization”)
 $Y \approx 0$ is an equality involving variable Y with coefficient 1
Resolution of the equality against Y
 $\mathcal{S} \vdash Y \approx 0$ (already proved)
Introduction of the value of Y into $Y \preceq -1$
 $\mathcal{S} \vdash 0 \preceq -1$ (proved by “eq as ineq”)
 $\mathcal{S} \vdash 0 \preceq -1$ Is a constraint without any variables.
 $\mathcal{S} \vdash 0 \leq -1$ (proved by “leq on \mathbb{Z} ”)
 $\mathcal{S} \vdash \perp$

Figure 4: Example of a resolution using Omega2FOL (outline)

$$\begin{array}{c}
 \frac{\frac{\frac{\frac{\frac{\frac{\frac{}{\vdash X \succeq y}}{\vdash X \ominus Y \succeq 0}}{\text{geqgeq}}}{\vdash X \succ y}}{\vdash X \ominus Y \ominus 1 \succeq 0}}{\text{gtgeq}}}{\vdash Y \ominus X \succeq 0}}{\text{leqgeq}}}{\vdash Y \ominus X \ominus 1 \succeq 0}}{\text{ltgeq}} \\
 \frac{\frac{\frac{\frac{\frac{\frac{\frac{}{\vdash X \approx Y}}{\vdash X \succeq Y}}{\text{eqgeq}}}{\vdash 0 \succeq 0}}{\text{geq0}}}{\vdash X \oplus n \otimes Y \succeq 0}}{\text{geqadd}}}{\vdash X \succeq 0 \quad \vdash Y \succeq 0 \quad m \otimes X \approx n \otimes Y \quad m \geq 0 \quad -1 - n \geq 0}}{\vdash \perp} \text{falsei} \\
 \frac{\vdash \perp}{\vdash a \otimes (X \ominus Y) \approx \ominus Z \quad \vdash Z \succeq 0} \\
 \frac{\vdash a \otimes (Y \ominus X) \approx \ominus T \quad \vdash T \succeq 0}{\vdash a \geq 0 \quad \vdash b \geq 0} \\
 \frac{\vdash a \geq 0 \quad \vdash b \geq 0}{\vdash X \approx Y} \text{eqi}
 \end{array}$$

Figure 5: Simplex-logic rules

Corollary :

Construct 3 *Solver2FOL's logic is complete with respect to linear integer constraints.*

The proof is straightforward : Solver2FOL's logic can express Simplex2FOL's results, Simplex2FOL can produce results for the same set of constraints as the Simplex, Simplex is complete with respect to linear integer constraints.

In order to build Solver2FOL proofs using the Simplex, the only things one has to do is :

- keep the core of the Simplex decision procedure
- modify the proof-generating components so as to compose proofs using either *modus ponens* on Solver2FOL lemmas or lemma composition of Solver2FOL lemmas instead of rules composition of Simplex-logic rules. The example of such a modification is given on figure 6.

This transformation procedure is straightforward - and probably automatizable, once the lemmas have been proved. As a matter of fact, it does not require understanding the algorithm itself, only changing its return statements.

6 Properties of Solver2FOL

6.1 Representation power

6.1.1 Introduction

It is clear that any set of inequalities and equalities involving only integer linear relations can be represented using Solver2FOL. It is also clear that polynomial constraints can be represented using Solver2FOL.

A valid question is : can more complex conditions be represented using Solver2FOL ?

6.1.2 Expressing complex functions through brute force

A function such as $n \mapsto \lfloor 2^n \rfloor$ cannot be represented as a finite expression in Solver2FOL's expression language on \mathbb{N} .

However, for any value of N , as figure 7 shows, it can be represented as one disjunction of $\mathcal{O}(2^N)$ conjunctions of equalities on $] - 2^N, 2^N]$ plus one inequality. If we give to N a good

Simplex-logic style : P, Q are proofs, r is the number of current row, $maxProof$ is an auxiliary function, $geqct(c)$ is a construct meaning “ c is non-negative”

```

mkEqProof( $X, Y$ ) =
   $r \leftarrow r + 1$ 
  fill row  $r$  with coefficients for  $X \ominus Y$ 
  ( $a, Z, P$ )  $\leftarrow maxProof(r)$ 
  fill row  $r$  with coefficients for  $Y \ominus X$ 
  ( $b, T, Q$ )  $\leftarrow maxProof(r)$ 
   $r \leftarrow r - 1$ 
  return eqi(  $arith(a \otimes (X \ominus Y), \ominus Z), P, geqct(a),$ 
                 $arith(b \otimes (Y \ominus X), \ominus T), Q, geqct(b)$  )

```

Solver2FOL style : the return statement changes and becomes

```

return lemma “equi”[“a”  $\leftarrow a$ , “b”  $\leftarrow b$ 
                 “X”  $\leftarrow X$ , “Y”  $\leftarrow Y$ , “Z”  $\leftarrow Z$ 
                 “T”  $\leftarrow T$ ]( $P, Q$ )

```

Figure 6: Translation from Simplex-logic style to Solver2FOL (fragment)

value, say the size in bits of an memory word, we have, as far as the computer is concerned, represented our function.

As a matter of fact, all functions can be represented with Solver2FOL using such a trick, provided they stay into the integer domain of the computer. However, since this form potentially requires as many disjunctions as there are integer which can be represented by the

$$x \mapsto \lfloor f(x) = 2^x \rfloor$$

or

$$\bigvee_{i=0}^{i=2^N} (x \approx i \wedge f(x) \approx 2^i) \vee (x < 0 \wedge f(x) \approx 0)$$

Figure 7: A represented non-representable function.

machines, or as there are memory words which can be addressed by the hardware, this simple remark does not mean much about the expressiveness of the language : such representations are, computationally speaking, infinite.

6.1.3 Lessons from the brute-force approach

The brute force approach teaches us that the real limitation of the expressiveness using Solver2FOL is in fact a size limitation.

6.1.3.1 Disjunctive Normal Forms

Property 1 *Let us consider $t : k \mapsto 2^k$. Let $[0, B[$ be the range of addressable memory in a machine M . Let S be the memory size of an representation of t in Disjunctive Normal Form (DNF) using Solver2FOL for array bounds checking elimination or dynamic memory protection elimination on M .*

$$S > B$$

Corollary : As such, t cannot be represented in DNF using Solver2FOL.

Proof of property 1 *Let $]-R, R]$ be the range of representable integers on a given computer.*

The simple study conducted in annex B.1 proves that any representation of function t in Disjunctive Normal Form using Solver2FOL will require at least $R/4$ terms.

If we assume that Solver2FOL is used for dynamic array bounds checking elimination or for dynamic memory checking elimination, we need the range of computable integers to be able to represent the whole range of addressable memory. In other words, $R \geq B$.

If we assume that each term will require at least eight bytes of memory allocation, the total amount of memory required for representation of t will be at least $S = 2 \cdot R$.

Hence,

$$S > B$$

6.1.3.2 Recursivity However, as shown in figure 8 a logic with some notion of recursive definition would allow a simple definition of function t in DNF. Such a logic would allow a more complete or/and simpler expression language.

$$T(x, y) \equiv_{def} \begin{cases} x \prec 0 \wedge y \approx 0 \\ \vee x \approx 0 \wedge y \approx 1 \\ \vee T(x \ominus 1, z) \wedge z \approx y \setminus 2 \end{cases}$$

Figure 8: A possible definition of $t : n \mapsto 2^n$

6.1.4 Summary

Solver2FOL’s expression language is powerful enough to represent linear and polynomial expressions. Since most operations which require integer constraints solving, such as array bounds checking optimization of automatic loops parallelization, most often imply only linear constraints, seldom polynomials, and almost never more complicated expressions, this language can be considered complete enough.

However, more specialized security policies might require even more expressive languages. We handle this issue in section 7.2.

6.2 Expressiveness of proofs

As proved in section 5, Solver2FOL is complete with respect to linear integer constraints.

What can be said beyond that ?

Well, we can say that Solver2FOL knows no such thing as an induction proof. This means that theorems handling generalized sums, such as lemma 7, cannot be expressed entirely within Solver2FOL’s logic. *They must rely on a meta-logic and, for each given sum met during a proof, they must be instantiated for the number of terms in the sum.*

This means that Solver2FOL will produce numerous large proofs where there could have been only one relatively simple proof in a proof library. This issue is also handled in section 7.2 .

7 From theory to implementation

7.1 Integrating Solver2FOL in a real system

So far, Solver2FOL has not been introduced as a part of a real system. We plan to incorporate this work into the FLINT system [15] by extending the typed intermediate language with dependent types. In the following, we briefly discuss how Solver2FOL may be integrated with

some existing dependently typed languages.

Xi’s dependent type system [17], which has been used for array-bounds checking elimination, relies on an external constraints solver, symbolized by \models in Xi [20]. In practice, for integers, this constraints solver is some (simplified) variant of the Omega test.

Since we are replacing the Omega test by Omega2FOL, one of the next natural steps would be to insert Omega2FOL in the dependent type system. However, this is not as trivial as it sounds, since an implicit requirement on the external constraints solver seems to be completion. A lack of completion resulting in an incomplete type-checker and, as far as PCC is concerned, the rejection of perfectly sound programs for arbitrary reasons. Of course, the Omega test is not complete on the set of diophantine equations. No algorithm is.

So, how do we introduce Solver2FOL in here ? A convenient answer would be that if Solver2FOL can do everything the Omega test can, then of course it can serve as the external solver in a dependent type system. However, this is not true. Since the external solver is supposed complete, neither the Omega test nor Omega2FOL nor any other algorithm is powerful enough as to serve as such a solver.

Before introducing effectively Omega2FOL in the dependent type system, we will have to determine exactly what set of constraints can be solved with Omega2FOL or, for that matter, any *2FOL solver.

Or we can use another approach. After all, the whole point of Solver2FOL is not to be locked with any solver algorithm, even if it is a *2FOL solver. Turning the dependent type system from Omega test-dependent to Omega2FOL-dependent is only a small improvement.

The idea would be, simply put, to make the dependent type system solver-independent. In other words, replacing \models by \vdash . Replacing “my algorithm can prove this” by “there is an algorithm which can prove this, here is how”. And this algorithm may be Omega2FOL, Simplex2FOL, *2FOL. Or Twelf [12], or Coq [4], or the user himself. And the algorithm does not have to be trusted.

This is one of the next steps in the evolution of Solver2FOL.

7.2 Twelf

We have begun working on integration using the Edinburgh Logical Framework (LF) [2] through Twelf. In other words, we are trying to encode the whole Solver2FOL system into LF and to translate proofs into the Twelf system. This task has been undertaken with the support of Princeton’s *PCC team*, using PCC team’s meta-logic for Twelf.

Using some meta-language (such as LF) and some concrete implementation of it (like Twelf)

to describe Solver2FOL allows to turn Solver2FOL into a really trustable and extensible part of the trusted computing base. Extensions to Solver2FOL are new definitions, such as new operators, and new hard-to-prove-but-fully-trusted “axioms”. For example, Fermat’s theorem might be needed some day or maybe some specialized theorems related to bank accounting.

Additionally, the Twelf meta-logic we are using allows us a reasoning of somewhat higher order. Axioms of induction on \mathbb{N} , for example, let us overcome the size problem of induction proofs (cf. section 6.2). New functions can be defined and seamlessly integrated, which is a solution to the potential expressiveness problem presented in section 6.1.4.

8 Future work

8.1 Implementation

Our implementation of Solver2FOL in Twelf is not complete yet, since the PCC team’s libraries we are basing our work on are still in early development. For example, since the libraries do not include full support for lists yet, and since Σ sums rely on lists, we have had to pause development to contribute to this non-trivial part of the libraries.

8.2 Extensions to non-linear constraints

Recent versions of the Omega test and of other tests handle some limited kinds of non-linear constraints, using several techniques such as polynomials factoring or linear approximation on segments. We are planning to prove as many as possible of these extensions using Solver2FOL and to add them to Omega2FOL.

8.3 Exceptions handling

As explained earlier, the semantics of Solver2FOL do not handle integer arithmetic exceptions in a fully satisfactory way. One satisfactory way would be to introduce some “non-bounded” integers, using either unary integers or some fancy Big Integer construct. The first possibility would be simpler, hence easier to trust.

Another way would be to create proof templates instead of proofs. In this case, the templates would be instantiated at check-time with platform-dependent information. We have not pursued this trail any further yet. It promises to be very complicated as well as very rewarding.

9 Related work

The idea of PCC was first proposed by Necula and Lee [10]. To handle integer constraints, they introduce the Simplex-Logic, a form of minimalistic logic tailored for proving results obtained by the Simplex algorithm. However, since this logic is very specific, they can neither express proofs using any other algorithm or extend their algorithm to more generic constraints. Xi [20, 18], on the other hand, proposes a dependent type system, which totally abstracts the the solver algorithm, considering it as a prerequisite. Several other works require integer constraints solving algorithms in PCC and would benefit from a generic framework. Among them, TAL [7], Crary and Weirich [5]’s resource bound certification, Wang and Appel [16]’s safe garbage collection and an attempt at making the whole PCC system more generic proposed by Appel and Felty [1].

10 Conclusion

We presented a feasibility study for the elaboration of an algorithm independent framework for verifying and proving integer constraints. Our framework, Solver2FOL, is simple and can represent all commonly used integer constraints. Moreover, although its proof power is necessarily limited, it can be used at least on the whole range of linear integer constraints, which are by far the most common constraints encountered at compilation-time. We also presented our solution for overcoming expressiveness limitations and overgrown proofs size for its implementation in Twelf. We plan to implement this framework into the FLINT system [15].

References

- [1] A. Appel and A. Felty. A semantic model of types and machine instructions for proof-carrying code. In *Proc. 27th Annual ACM SIGPLAN-SIGACT Symp. on Principles of Programming Languages*, pages 243–253. ACM Press, 2000.
- [2] A. Avron, F. Honsell, and I. Mason. An overview of the edinburgh logical framework, 1989.
- [3] W. W. Bledsoe. The Sup-Inf method in praesburger arithmetic. Technical report, University of Texas Math Dept., December 1974.
- [4] C. Cornes. Compilation du filtrage de motifs avec types dépendants dans le système coq. In *Actes des Journées du GDR Programmation 1996*, Orleans, France, Novembre 1996.
- [5] K. Crary and S. Weirich. Resource bound certification. In *Proc. 27th Annual ACM SIGPLAN-SIGACT Symp. on Principles of Programming Languages*. ACM Press, 2000.

- [6] G. B. Dantzig and B. C. Eaves. Fourier-Motzkin elimination and its dual. *J. Combin. Theo. A*, 14:288–297, 1973.
- [7] G. Morrisett, D. Walker, K. Crary, and N. Glew. From system F to typed assembly language. In *Proc. 25rd Annual ACM SIGPLAN-SIGACT Symp. on Principles of Programming Languages*, pages 85–97. ACM Press, 1998.
- [8] G. Necula. Proof-carrying code. In *Twenty-Fourth Annual ACM Symp. on Principles of Prog. Languages*, pages 106–119, New York, Jan 1997. ACM Press.
- [9] G. Necula and P. Lee. Safe, untrusted agents using proof-carrying code. In *Special Issue on Mobile Agent Security: LNCS Vol 1419*. Springer-Verlag, October 1997.
- [10] G. C. Necula. *Compiling with Proofs*. PhD thesis, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, Sept. 1998.
- [11] G. Nelson. Techniques for program verification. Technical Report CSL-81-10, Xerox Palo Alto Research Center, 1981.
- [12] F. Pfenning and C. Schürmann. System description: Twelf — a meta-logical framework for deductive systems. In H. Ganzinger, editor, *Proceedings of the 16th International Conference on Automated Deduction (CADE-16)*, pages 202–206, Trento, Italy, July 1999. Springer-Verlag LNAI 1632.
- [13] W. Pugh. A practical algorithm for exact array dependence analysis. *Commun. ACM*, 35(8):102–115, August 1992.
- [14] W. Pugh and D. Wonnacott. An exact method for analysis of value-based array data dependences. In *Proceedings of the Sixth Annual Workshop on Programming Languages and Compilers for Parallel Computing*, Dec 93.
- [15] Z. Shao. An overview of the FLINT/ML compiler. In *Proc. 1997 ACM SIGPLAN Workshop on Types in Compilation*. Published as Boston College Computer Science Dept. Technical Report BCCS-97-03, June 1997.
- [16] D. Wang and A. Appel. Safe garbage collection = regions + intentional type analysis. Technical report, Dept. of Computer Science, Princeton University, July 1999.
- [17] H. Xi. *Dependent Types in Practical Programming*. PhD thesis, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, Sept. 1998.
- [18] H. Xi and R. Harper. A dependently typed assembly language. Technical Report CMU-CS-99-xxx, School of Computer Science, Carnegie Mellon University, Pittsburgh, PA, July 1999.
- [19] H. Xi and F. Pfenning. Eliminating array bound checking through dependent types. In *Proc. ACM SIGPLAN '98 Conf. on Prog. Lang. Design and Implementation*, pages 249–257, New York, 1998. ACM Press.

- [20] H. Xi and F. Pfenning. Dependent types in practical programming. In *Proceedings of ACM SIGPLAN Symposium on Principles of Programming Languages*, pages 214–227, San Antonio, January 1999.

A Proofs using Solver2FOL

A.1 Foreword

This appendix contains purely formal proofs built using Solver2FOL.

A.2 Syntactical conventions

- A name or a number alone in a formal proof is a reference to the corresponding lemma, theorem, axiom, definition. Only sublemmas are referred to by their number. Other lemmas, theorems, axioms, definitions are referred to by their name.

Using somelemma

- Notation $\frac{}{H \vdash \blacksquare}$ is a shortcut for the utilization of the *modus ponens* on an instantiation of lemma (or theorem, or axiom, ...) *sommelemma*.
- $\Sigma_{i=1}^n X_i$ is a shortcut for $((X_1 \oplus X_2) \oplus \dots X_n)$
- $\Sigma_{i \in I} X_i$ is a shortcut for some left-parenthesized sum (using \oplus) of all terms of $(X_i)_{i \in I}$ ¹

A.3 Basic theorems

Proof of basic theorem 1 Let $H \equiv_{def} 0 \leq e \wedge e < b \wedge a \approx b \otimes c \oplus e$ and $K \equiv_{def} 0 \leq d \wedge d < b \wedge a = b \cdot c + d$.

Sub lemma 1 $\vdash b \otimes c \oplus e \approx b \cdot c \oplus e$

Proof of sub lemma 1

$$\frac{\frac{\text{Using "mult on } \mathbb{Z}"}{\vdash b \otimes c \approx b \cdot c} \quad \frac{\text{Using "add eq"}}{\vdash \blacksquare}}{\vdash b \otimes c \oplus e \approx b \cdot c \oplus e}}$$

¹The left-parenthesized convention has been adopted for the sake of simplicity of proofs. However, there is no obligation of expressing sums like this.

Sub lemma 2 $H \vdash a \approx b \cdot c \oplus e$

Proof of sub lemma 2

$$\frac{\frac{H \vdash a \approx b \otimes c \oplus e \quad \frac{\text{Using 1}}{\vdash b \otimes c \oplus e \approx b \cdot c \oplus e}}{H \vdash a \approx b \otimes c \oplus e \wedge b \otimes c \oplus e \approx b \cdot c \oplus e} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash a \approx b \cdot c \oplus e}$$

Sub lemma 3 $H \vdash a = b \cdot c + e$

Proof of sub lemma 3

$$\frac{\frac{\frac{\text{Using 2}}{H \vdash a \approx b \cdot c \oplus e} \quad \frac{\text{Using "add on } \mathbb{Z}}{H \vdash b \cdot c \oplus e \approx b \cdot c + e}}{H \vdash a \approx b \cdot c \oplus e \wedge b \cdot c \oplus e \approx b \cdot c + e} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{\frac{H \vdash a \approx b \cdot c + e}{H \vdash a = b \cdot c + e}} \quad \frac{\text{Using "eq on } \mathbb{Z}}{H \vdash \blacksquare}}$$

Sub lemma 4 $a \setminus b \approx c \vdash \lfloor \frac{a}{b} \rfloor = c$

Proof of sub lemma 4

$$\frac{\frac{\frac{H \vdash 0 \leq e \quad H \vdash e < b \quad \frac{\text{Using 3}}{H \vdash a = b \cdot c + e}}{H \vdash 0 \leq e \wedge e < b \wedge a = b \cdot c + e}}{\exists e, 0 \leq e \wedge e < b \wedge a \approx b \otimes c \oplus e \vdash \exists e, 0 \leq e \wedge e < b \wedge a = b \cdot c + e}}{a \setminus b \approx c \vdash \lfloor \frac{a}{b} \rfloor = c}$$

Sub lemma 5 $\vdash b \cdot c + d \approx b \otimes c \oplus d$

Proof of sub lemma 5

$$\frac{\frac{\frac{\text{Using 1}}{\vdash b \otimes c \oplus d \approx b \cdot c \oplus d} \quad \frac{\text{Using "add in } \mathbb{Z}}{\vdash b \cdot c \oplus d \approx b \cdot c + d}}{\vdash b \otimes c \oplus d \approx b \cdot c \oplus d \wedge b \cdot c \oplus d \approx b \cdot c + d} \quad \frac{\text{Using "eq ref"}}{\vdash \blacksquare}}{\frac{\vdash b \otimes c \oplus d \approx b \cdot c + d}{\vdash b \cdot c + d \approx b \otimes c \oplus d}}$$

Sub lemma 6 $K \vdash a \approx b \otimes c \oplus d$

Proof of sub lemma 6

$$\frac{\frac{K \vdash a = b \cdot c + d}{K \vdash a \approx b \cdot c + d} \quad \frac{\text{Using 5}}{K \vdash b \cdot c + d \approx b \otimes c \oplus d}}{\frac{K \vdash a \approx b \cdot c + d \wedge b \cdot c + d \approx b \otimes c \oplus d}{K \vdash a \approx b \otimes c \oplus d}} \quad \frac{\text{Using "eq trans"}}{K \vdash \blacksquare}$$

Sub lemma 7 $\lfloor \frac{a}{b} \rfloor = c \vdash a \setminus b \approx c$

Proof of sub lemma 7

$$\frac{\frac{\frac{K \vdash 0 \leq d \quad K \vdash d < b}{K \vdash 0 \leq d \wedge d < b} \quad \frac{\text{Using 6}}{K \vdash a \approx b \otimes c \oplus d}}{K \vdash 0 \leq d \wedge d < b \wedge a \approx b \otimes c \oplus d}}{\frac{\exists d, 0 \leq d \wedge d < b \wedge a = b \cdot c + d \vdash \exists d, 0 \leq d \wedge d < b \wedge a \approx b \otimes c \oplus d}{\lfloor \frac{a}{b} \rfloor = c \vdash a \setminus b \approx c}}$$

Hence,

$$\frac{\frac{\frac{\text{Using 7}}{\lfloor \frac{a}{b} \rfloor = c \vdash a \setminus b \approx c}}{\vdash \lfloor \frac{a}{b} \rfloor = c \Rightarrow a \setminus b \approx c} \quad \frac{\frac{\text{Using 4}}{a \setminus b \approx c \vdash \lfloor \frac{a}{b} \rfloor = c}}{\vdash a \setminus b \approx c \Rightarrow \lfloor \frac{a}{b} \rfloor = c}}{\vdash a \setminus b \approx c \Leftrightarrow \lfloor \frac{a}{b} \rfloor = c}}$$

Proof of basic theorem 2 Let $H \equiv_{def} 0 \leq e \wedge e < b \wedge a \approx b \otimes c \oplus e$ and $K \equiv_{def} 0 \leq e \wedge e < b \wedge a = b \cdot c + e$.

Sub lemma 8 $a \% b \approx e \vdash a \bmod b = e$

Proof of sub lemma 8

$$\frac{\frac{\frac{H \vdash 0 \leq e \quad H \vdash e < b}{H \vdash 0 \leq e \wedge e < b} \quad \frac{\text{Using 3}}{H \vdash a = b \cdot c + e}}{H \vdash 0 \leq e \wedge e < b \wedge a = b \cdot c + e}}{\frac{\exists c, 0 \leq e \wedge e < b \wedge a \approx b \otimes c \oplus e \vdash \exists d, 0 \leq e \wedge e < b \wedge a = b \cdot d + e}{a \% b \approx e \vdash a \bmod b = e}}$$

Sub lemma 9 $a \bmod b = e \vdash a \% b \approx e$

Proof of sub lemma 9

$$\frac{\frac{\frac{K \vdash 0 \leq e \quad K \vdash e < b \quad \frac{\text{Using 6}}{K \vdash a \approx b \otimes d \oplus e}}{K \vdash 0 \leq e \wedge e < b \wedge a \approx b \otimes d \oplus e}}{\exists c, 0 \leq e \wedge e < b \wedge a = b \cdot d + e \vdash \exists c, 0 \leq e \wedge e < b \wedge a \approx b \otimes c \oplus e}}{a \bmod b = e \vdash a \% b \approx e}$$

Hence,

$$\frac{\frac{\frac{\text{Using 8}}{a \% b \approx e \vdash a \bmod b = e} \quad \frac{\text{Using 9}}{a \bmod b = e \vdash a \% b \approx e}}{\vdash a \% b \approx e \Rightarrow a \bmod b = e \quad \vdash a \bmod b = e \Rightarrow a \% b \approx e}}{\vdash a \% b \approx e \Rightarrow a \bmod b = e \wedge a \bmod b = e \Rightarrow a \% b \approx e}}{\vdash a \% b \approx e \iff a \bmod b = e}$$

Proof of basic theorem 3

$$\frac{\text{Using "mod on } \mathbb{Z}\text{"}}{\frac{\vdash a \% b \approx 0 \iff a \bmod b = 0}{\vdash a \parallel b \iff a|b}}$$

Proof of basic theorem 4 let $H \equiv_{def} 0 \leq c \wedge a \approx b \oplus c$ and $K \equiv_{def} 0 \leq d \wedge a = b + d$

Sub lemma 10 $a \succeq b \vdash a \geq b$

Proof of sub lemma 10

$$\frac{\frac{\frac{H \vdash a \approx b \oplus c \quad \frac{\text{Using "add in } \mathbb{Z}\text{"}}{H \vdash b \oplus c \approx b + c}}{H \vdash a \approx b \oplus c \wedge b \oplus c \approx b + c} \quad \frac{\text{Using "eq in } \mathbb{Z}\text{"}}{H \vdash \blacksquare}}{H \vdash a \approx b + c} \quad \frac{H \vdash 0 \leq c}{H \vdash a = b + c}}{\frac{H \vdash 0 \leq c \wedge a = b + c}{\exists c, 0 \leq c \wedge a \approx b \oplus c \vdash \exists d, 0 \leq d \wedge a = b + d}}{a \succeq b \vdash a \geq b}$$

Sub lemma 11 $a \geq b \vdash a \succeq b$

Proof of sub lemma 11

$$\frac{\frac{\frac{K \vdash a = b + d \quad \text{Using "add in } \mathbb{Z}\text{"}}{K \vdash a \approx b + d} \quad \frac{K \vdash b + d \approx d}{\text{Using "eq trans"}}$$

$$\frac{\frac{K \vdash a \approx b + d \wedge b + d \approx d}{K \vdash \blacksquare}}{K \vdash a \approx b \oplus d} \quad \frac{K \vdash 0 \leq d}{\exists d, 0 \leq d \wedge a = b + d \vdash \exists c, 0 \leq c \wedge a \approx b \oplus c}}{a \geq b \vdash a \succeq b}$$

Hence,

$$\frac{\frac{\text{Using 10}}{a \succeq b \vdash a \geq b} \quad \frac{\text{Using 11}}{a \geq b \vdash a \succeq b}}{\vdash a \succeq b \Rightarrow a \geq b} \quad \frac{\text{Using 11}}{a \geq b \vdash a \succeq b}}{\vdash a \geq b \Rightarrow a \succeq b}}{\vdash a \succeq b \Rightarrow a \geq b \wedge a \geq b \Rightarrow a \succeq b}}{\vdash a \succeq b \iff a \geq b}}$$

A.3.0.1 Opposite addition

Proof of main lemma 1

$$\frac{\frac{\text{Using "eq mult"}}{\vdash Y \ominus Y \approx 0 \otimes Y} \quad \frac{\text{Using "0 mult"}}{\vdash 0 \otimes Y \approx 0} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{\vdash Y \ominus Y \approx 0}}$$

A.3.0.2 "Distribution" of addition on equality 2

Proof of main lemma 2 Sub lemma 12 $X \approx Y \vdash X \oplus Z \approx Z \oplus Y$

Proof of sub lemma 12

$$\frac{\frac{\text{Using "add eq"}}{X \approx Y \vdash X \oplus Z \approx Y \oplus Z} \quad \frac{\text{Using "add comm"}}{\vdash Z \oplus Y \approx Y \oplus Z}}{X \approx Y \vdash X \oplus Z \approx Y \oplus Z \wedge Z \oplus Y \approx Y \oplus Z}}{\frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{X \approx Y \vdash X \oplus Z \approx Z \oplus Y}}$$

Hence,

$$\frac{\frac{\text{Using 12}}{\vdash Z \oplus X \approx X \oplus Z} \quad \frac{\text{Using "add comm"}}{\vdash Z \oplus X \approx X \oplus Z}}{X \approx Y \vdash X \oplus Z \approx Z \oplus Y \wedge Z \oplus X \approx X \oplus Z}}{\frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{X \approx Y \vdash Z \oplus X \approx Z \oplus Y}}$$

A.3.0.3 "Distribution" of multiplication on equality 2

Proof of main lemma 3 Sub lemma 13 $X \approx Y \vdash X \otimes Z \approx Z \otimes Y$

Proof of sub lemma 13

$$\frac{\frac{\text{Using "mult eq"}}{X \approx Y \vdash X \otimes Z \approx Y \otimes Z} \quad \frac{\text{Using "mult comm"}}{\vdash Z \otimes Y \approx Y \otimes Z}}{X \approx Y \vdash X \otimes Z \approx Y \otimes Z \wedge Z \otimes Y \approx Y \otimes Z}}{\frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{X \approx Y \vdash X \otimes Z \approx Z \otimes Y}}$$

Hence,

$$\frac{\frac{\text{Using } 13}{\vdash Z \otimes X \approx X \otimes Z} \quad \frac{\text{Using "mult comm"}}{\vdash Z \otimes X \approx X \otimes Z}}{X \approx Y \vdash X \otimes Z \approx Z \otimes Y \wedge Z \otimes X \approx X \otimes Z} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{X \approx Y \vdash Z \otimes X \approx Z \otimes Y}$$

Proof of basic theorem 5 Reflexivity :

$$\frac{\frac{\text{Using "0 add"}}{\vdash X \oplus 0 \approx X} \quad \frac{\text{Using "eq ref"}}{\vdash \blacksquare}}{\vdash 0 \leq 0 \quad \vdash X \approx X \oplus 0}}{\frac{\vdash 0 \leq 0 \wedge X \approx X \oplus 0}}{\vdash \exists i, 0 \leq i \wedge X \approx X \oplus i}}{\vdash X \succeq X}$$

Antisymmetry : let $H \equiv_{def} 0 \leq a \wedge 0 \leq b \wedge X \approx Y \oplus a \wedge Y \approx X \oplus b$

Sub lemma 14 $H \vdash Y \oplus a \approx X$

Proof of sub lemma 14

$$\frac{\frac{\text{Using "add trans"}}{H \vdash Y \oplus a \approx a \oplus Y} \quad H \vdash a \oplus Y \approx X \quad \text{Using "eq trans"}}{H \vdash Y \oplus a \approx a \oplus Y \wedge a \oplus Y \approx X} \quad H \vdash \blacksquare}{H \vdash Y \oplus a \approx X}$$

Sub lemma 15 $H \vdash X \oplus b \oplus a \approx Y \oplus a$

Proof of sub lemma 15

$$\frac{H \vdash Y \approx X \oplus b \quad \frac{\text{Using "eq ref"}}{H \vdash \blacksquare}}{H \vdash X \oplus b \approx Y} \quad \frac{\text{Using "add eq"}}{H \vdash \blacksquare}}{H \vdash X \oplus b \oplus a \approx Y \oplus a}$$

Sub lemma 16 $H \vdash X \oplus b \oplus a \approx X$

Proof of sub lemma 16

$$\frac{\frac{\text{Using } 15}{H \vdash X \oplus b \oplus a \approx Y \oplus a} \quad \frac{\text{Using } 14}{H \vdash Y \oplus a \approx X}}{H \vdash X \oplus b \oplus a \approx Y \oplus a \wedge Y \oplus a \approx X} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash X \oplus b \oplus a \approx X}$$

Sub lemma 17 $H \vdash X \oplus (b \oplus a) \approx X$

Proof of sub lemma 17

$$\frac{\frac{\text{Using 16}}{H \vdash X \oplus b \oplus a \approx X} \quad \frac{\text{Using "add assoc"}}{H \vdash X \oplus (b \oplus a) \approx X \oplus b \oplus a}}{H \vdash X \oplus b \oplus a \approx X \wedge X \oplus (b \oplus a) \approx X \oplus b \oplus a} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash X \oplus (b \oplus a) \approx X}$$

Sub lemma 18 $H \vdash (b \oplus a) \oplus X \approx X$

Proof of sub lemma 18

$$\frac{\frac{\text{Using 17}}{H \vdash X \oplus (b \oplus a) \approx X} \quad \frac{\text{Using "add comm"}}{H \vdash (b \oplus a) \oplus X \approx X \oplus (b \oplus a)}}{H \vdash X \oplus (b \oplus a) \approx X \wedge (b \oplus a) \oplus X \approx X \oplus (b \oplus a)} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash (b \oplus a) \oplus X \approx X}$$

Sub lemma 19 $H \vdash (b + a) \oplus X \approx (a \oplus b) \oplus X$

Proof of sub lemma 19

$$\frac{\frac{\text{Using "add on } \mathbb{Z}}{H \vdash (b + a) \approx (b \oplus a)} \quad \frac{\text{Using "add eq"}}{\vdash \blacksquare}}{H \vdash (b + a) \oplus X \approx (b \oplus a) \oplus X}}$$

Sub lemma 20 $H \vdash (b + a) \oplus X \ominus X \approx X \ominus X$

Proof of sub lemma 20

$$\frac{\frac{\frac{\text{Using 18}}{H \vdash (b \oplus a) \oplus X \approx X} \quad \frac{\text{Using 19}}{H \vdash (b + a) \oplus X \approx (b \oplus a) \oplus X}}{H \vdash (b \oplus a) \oplus X \approx X \wedge (b + a) \oplus X \approx (b \oplus a) \oplus X} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash (b + a) \oplus X \approx X} \quad \frac{\text{Using "add eq"}}{H \vdash \blacksquare}}{H \vdash (b + a) \oplus X \ominus X \approx X \ominus X}$$

Sub lemma 21 $\vdash (b + a) \oplus (X \ominus X) \approx (b + a) \oplus 0$

Proof of sub lemma 21

$$\frac{\frac{\text{Using "opp add"}}{\vdash (X \ominus X) \approx 0} \quad \frac{\text{Using "add eq 2"}}{\vdash \blacksquare}}{\vdash (b + a) \oplus (X \ominus X) \approx (b + a) \oplus 0}}$$

Sub lemma 22 $H \vdash (b + a) \approx (b + a) \oplus (X \ominus X)$

Proof of sub lemma 22

$$\frac{\frac{\frac{\text{Using 21}}{\vdash (b + a) \oplus (X \ominus X) \approx (b + a) \oplus 0} \quad \frac{\text{Using "0 add"}}{\vdash (b + a) \oplus 0 \approx b + a}}{\vdash (b + a) \oplus (X \ominus X) \approx (b + a) \oplus 0 \wedge (b + a) \oplus 0 \approx b + a}}{\vdash (b + a) \oplus (X \ominus X) \approx (b + a)} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{\vdash (b + a) \approx (b + a) \oplus (X \ominus X)} \quad \frac{\text{Using "eq ref"}}{\vdash \blacksquare}$$

Sub lemma 23 $H \vdash (b + a) \approx (b + a) \oplus X \ominus X$

Proof of sub lemma 23

$$\frac{\frac{\text{Using 22}}{H \vdash (b + a) \approx (b + a) \oplus (X \ominus X)} \quad \frac{\text{Using "add assoc"}}{H \vdash (b + a) \oplus (X \ominus X) \approx ((b + a) \oplus X) \ominus X}}{H \vdash (b + a) \approx (b + a) \oplus (X \ominus X) \wedge (b + a) \oplus (X \ominus X) \approx ((b + a) \oplus X) \ominus X}}{H \vdash (b + a) \approx ((b + a) \oplus X) \ominus X}$$

Sub lemma 24 $H \vdash a + b \approx X \ominus X$

Proof of sub lemma 24

$$\frac{\frac{\text{Using 20}}{H \vdash (b + a) \oplus X \ominus X \approx X \ominus X} \quad \frac{\text{Using 23}}{H \vdash (b + a) \approx (b + a) \oplus X \ominus X}}{H \vdash (b + a) \oplus X \ominus X \approx X \ominus X \wedge (b + a) \approx (b + a) \oplus X \ominus X}}{H \vdash a + b \approx X \ominus X}$$

Sub lemma 25 $H \vdash a \approx 0$

Proof of sub lemma 25

$$\frac{\frac{\frac{\text{Using 24}}{H \vdash a + b \approx X \ominus X} \quad \frac{\text{Using "opp add"}}{H \vdash X \ominus X \approx 0}}{H \vdash a + b \approx X \ominus X \wedge X \ominus X \approx 0}}{H \vdash a + b \approx 0}}{H \vdash a + b = 0} \quad \frac{\text{Using "eq on } \mathbb{Z}}{H \vdash \blacksquare}}{H \vdash a \approx 0}$$

Sub lemma 26 $H \vdash a \oplus Y \approx Y \oplus 0$

Proof of sub lemma 26

$$\frac{\frac{\text{Using 25}}{H \vdash a \approx 0} \quad \frac{\text{Using "add eq"}}{H \vdash \blacksquare}}{H \vdash a \oplus Y \approx 0 \oplus Y} \quad \frac{\text{Using "add comm"}}{\vdash 0 \oplus Y \approx Y \oplus 0} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash a \oplus Y \approx 0 \oplus Y \wedge 0 \oplus Y \approx Y \oplus 0} \quad \frac{}{H \vdash a \oplus Y \approx Y \oplus 0}$$

Sub lemma 27 $H \vdash X \approx Y \oplus 0$

Proof of sub lemma 27

$$\frac{\frac{\text{Using "add comm"}}{H \vdash Y \oplus a \approx a \oplus Y} \quad \frac{\text{Using 26}}{H \vdash a \oplus Y \approx Y \oplus 0} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash Y \oplus a \approx a \oplus Y \wedge a \oplus Y \approx Y \oplus 0} \quad \frac{}{H \vdash \blacksquare}}{H \vdash X \approx Y \oplus a} \quad \frac{H \vdash Y \oplus a \approx Y \oplus 0}{H \vdash X \approx Y \oplus a \wedge Y \oplus a \approx Y \oplus 0} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash X \approx Y \oplus 0}$$

Hence,

$$\frac{\frac{\text{Using 27}}{H \vdash X \approx Y \oplus 0} \quad \frac{\text{Using "0 add"}}{\vdash Y \oplus 0 \approx Y} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash X \approx Y \oplus 0 \wedge Y \oplus 0 \approx Y} \quad \frac{}{H \vdash \blacksquare}}{\exists a, 0 \leq a \wedge X \approx Y \oplus a \wedge \exists b, 0 \leq b \wedge Y \approx X \oplus b \vdash X \approx Y} \quad \frac{X \succeq Y \wedge Y \succeq X \vdash X \approx Y}{\vdash X \succeq Y \wedge Y \succeq X \Rightarrow X \approx Y}$$

Transitivity : let $H \equiv_{def} 0 \leq a \wedge 0 \leq b \wedge X \approx Y \oplus a \wedge Y \approx Z \oplus b$

Sub lemma 28 $H \vdash Y \oplus a \approx Z \oplus b \oplus a$

Proof of sub lemma 28

$$\frac{\text{Using "add eq"}}{H \vdash Y \approx Z \oplus b} \quad \frac{}{H \vdash \blacksquare}}{H \vdash Y \oplus a \approx Z \oplus b \oplus a}$$

Sub lemma 29 $H \vdash X \approx Z \oplus b \oplus a$

Proof of sub lemma 29

$$\frac{\text{Using 28}}{H \vdash Y \oplus a \approx Z \oplus b \oplus a} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{H \vdash X \approx Y \oplus a \wedge Y \oplus a \approx Z \oplus b \oplus a} \quad \frac{}{H \vdash \blacksquare}}{H \vdash X \approx Z \oplus b \oplus a}$$

Sub lemma 30 $\vdash Z \oplus b \oplus a \approx Z \oplus (a + b)$

Proof of sub lemma 30

$$\frac{\frac{\text{Using "add assoc"}}{\vdash Z \oplus a \oplus b \approx Z \oplus (a \oplus b)} \quad \frac{\frac{\text{Using "add in } \mathbb{Z}}{\vdash (a \oplus b) \approx (a + b)} \quad \frac{\text{Using "eq/add dist2"}}{\vdash \blacksquare}}{\vdash Z \oplus (a \oplus b) \approx Z \oplus (a + b)}}{\vdash Z \oplus a \oplus b \approx Z \oplus (a \oplus b) \wedge Z \oplus (a \oplus b) \approx Z \oplus (a + b)} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{\vdash Z \oplus a \oplus b \approx Z \oplus (a + b)}$$

Sub lemma 31 $H \vdash X \approx Z \oplus (a + b)$

Proof of sub lemma 31

$$\frac{\frac{\text{Using 29}}{H \vdash X \approx Z \oplus a \oplus b} \quad \frac{\text{Using 30}}{H \vdash Z \oplus a \oplus b \approx Z \oplus (a + b)} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{\frac{H \vdash X \approx Z \oplus a \oplus b \wedge Z \oplus a \oplus b \approx Z \oplus (a + b)}{H \vdash X \approx Z \oplus (a + b)}}$$

Hence,

$$\frac{\frac{\text{Using 31}}{H \vdash 0 \leq (a + b) \wedge H \vdash X \approx Z \oplus (a + b)}}{H \vdash 0 \leq (a + b) \wedge X \approx Z \oplus (a + b)}}{H \vdash \exists c, 0 \leq c \wedge X \approx Z \oplus c} \quad \frac{X \succeq Y \wedge Y \succeq Z \vdash X \succeq Z}{X \succeq Y \wedge Y \succeq Z \vdash X \succeq Z}$$

Proof of basic theorem 6 *The proof is very similar to that of theorem 4*

Proof of basic theorem 7 *The proof is very similar to that of theorem 4*

Proof of basic theorem 8 *The proof is very similar to that of theorem 4*

Proof of basic theorem 9 *The proof is very similar to that of theorem 5*

Proof of basic theorem 10 **Sub lemma 32** $X \preceq Y \vdash Y \succeq X$

Proof of sub lemma 32

$$\frac{\exists a, 0 \leq a \wedge Y \approx X \oplus a \vdash \exists b, 0 \leq b \wedge Y \approx X \oplus b}{X \preceq Y \vdash Y \succeq X}$$

Sub lemma 33 $X \preceq Y \vdash Y \succeq X$

Proof of sub lemma 33

$$\frac{\exists a, 0 \leq a \wedge X \approx Y \oplus a \vdash \exists b, 0 \leq b \wedge X \approx Y \oplus b}{X \succeq Y \vdash Y \preceq X}$$

Hence,

$$\frac{\frac{\text{Using 32}}{X \preceq Y \vdash Y \succeq X} \quad \frac{\text{Using 33}}{X \succeq Y \vdash Y \preceq X}}{\vdash X \preceq Y \Rightarrow Y \succeq X \quad \vdash X \succeq Y \Rightarrow Y \preceq X}}{\vdash X \preceq Y \Rightarrow Y \succeq X \wedge X \succeq Y \Rightarrow Y \preceq X}}{\vdash X \preceq Y \iff Y \succeq X}$$

Proof of basic theorem 11 *The proof is very similar to that of theorem 10*

Proof of basic theorem 12 *(Note : This theorem/lemma is not “static”. cf. 6.2 page 27)*

Let $H \equiv_{def} 0 < a \wedge Y \approx X \oplus a$, $K \equiv_{def} x \approx$ and $L \equiv_{def} 0 \leq a \wedge Y \approx X \oplus a \wedge \neg(Y \approx A)$

Sub lemma 34 $L \vdash X \approx Y$

Proof of sub lemma 34

$$\frac{\frac{L \vdash Y \approx X \oplus 0 \quad \frac{\text{Using “eq ref”}}{L \vdash \blacksquare}}{L \vdash X \oplus 0 \approx Y} \quad \frac{\text{Using “0 add”}}{\vdash X \approx X \oplus 0}}{L \vdash X \oplus 0 \approx Y \wedge X \approx X \oplus 0}}{L \vdash X \approx Y}$$

Sub lemma 35 $L \vdash 0 < a$

Proof of sub lemma 35

$$\frac{\frac{L \vdash 0 < a}{L \vdash 0 \leq a} \quad \frac{\frac{\text{Using 34}}{L, 0 = a \vdash \neg(X \approx Y)} \quad L, 0 = a \vdash X \approx Y}{L, 0 = a \vdash X \approx Y \wedge \neg(X \approx Y)}}{\frac{L, 0 = a \vdash \perp}{L \vdash 0 \neq a}}}{L \vdash 0 \leq a \wedge 0 \neq a}}{L \vdash 0 < a}$$

Sub lemma 36 $X \preceq Y \vdash X \prec Y \vee X \approx Y$

Proof of sub lemma 36

$$\frac{\frac{\frac{\text{Using 35}}{L \vdash 0 < a} \quad L \vdash Y \approx X \oplus a}{L \vdash 0 < a \wedge Y \approx X \oplus a}}{L \vdash \exists b, 0 < b \wedge Y \approx X \oplus b}}{X \preceq Y \wedge \neg(X \approx Y) \vdash X \prec Y}}{X \preceq Y \vdash X \prec Y \vee X \approx Y}$$

Sub lemma 37 $K \vdash X \preceq Y$

Proof of sub lemma 37

$$\frac{\frac{\frac{\text{Using "0 add"} \quad \text{Using "eq ref"}}{\vdash X \approx X \oplus 0} \quad \vdash \blacksquare}}{K \vdash X \oplus 0 \approx X} \quad \frac{K \vdash X \approx Y \quad \text{Using "eq trans"}}{K \vdash \blacksquare}}{\frac{K \vdash X \oplus 0 \approx X \wedge X \approx Y}{K \vdash X \oplus 0 \approx Y} \quad \text{Using "eq ref"} \quad K \vdash \blacksquare}}{K \vdash 0 \leq 0} \quad \frac{K \vdash Y \approx X \oplus 0}{\frac{K \vdash 0 \leq 0 \wedge Y \approx X \oplus 0}{K \vdash \exists c, 0 \leq c \wedge Y \approx X \oplus i}}{K \vdash X \preceq Y}}$$

Sub lemma 38 $X \prec Y \vee X \approx Y \vdash X \preceq Y$

Proof of sub lemma 38

$$\frac{\frac{\frac{H \vdash 0 \leq a \quad H \vdash Y \approx X \oplus a}{H \vdash 0 \leq a \wedge Y \approx X \oplus a}}{\exists a, H, 0 \leq a \wedge Y \approx X \oplus a} \quad \text{Using 37}}{X \prec Y \vdash Y \preceq X} \quad \frac{\text{Using 37}}{K \vdash X \preceq Y}}{X \prec Y \vee K \vdash X \preceq Y}$$

Hence,

$$\frac{\frac{\text{Using 36}}{X \preceq Y \vdash X \prec Y \vee X \approx Y} \quad \frac{\text{Using 38}}{X \prec Y \vee X \approx Y \vdash X \preceq Y}}{\vdash X \preceq Y \Rightarrow (X \prec Y \vee X \approx Y) \vdash (X \prec Y \vee X \approx Y) \Rightarrow X \preceq Y}}{\vdash (X \preceq Y \Rightarrow (X \prec Y \vee X \approx Y)) \wedge ((X \prec Y \vee X \approx Y) \Rightarrow X \preceq Y)} \quad \vdash X \preceq Y \iff (X \prec Y \vee X \approx Y)$$

Proof of basic theorem 13 *The proof is very similar to that of theorem 12.*

Proof of basic theorem 14 *Let $H \equiv_{def} X \preceq Y \wedge X \succeq Y$*

$$\frac{H \vdash X \preceq Y \quad \frac{\text{Using "leq equiv geq"}}{H \vdash Y \preceq X}}{H \vdash X \preceq Y \wedge Y \preceq X} \quad \frac{\text{Using "lt antis"}}{H \vdash \blacksquare}}{X \preceq Y \wedge X \succeq Y \vdash X \approx Y}$$

Proof of basic theorem 15 *Let $H \equiv_{def} X \prec Y \wedge X \succ Y$*

$$\frac{\frac{\text{Using "lt as leq"}}{H \vdash X \preceq Y} \quad \frac{\text{Using "gt as geq"}}{H \vdash X \succeq Y}}{H \vdash X \preceq Y \wedge X \succeq Y} \quad \frac{\text{Using 34}}{H \vdash \neg(X \approx Y)}}{H \vdash X \approx Y \wedge \neg(X \approx Y)} \quad \frac{}{X \prec Y \wedge X \succ Y \vdash \perp}$$

A.3.0.4 Additive equality

Proof of main lemma 4 *Let $H \equiv_{def} X \approx Y \wedge Z \approx T$*

Sub lemma 39 $H \vdash X \oplus Z \approx X \oplus T$

Proof of sub lemma 39

$$\frac{H \vdash Z \approx T \quad \frac{\text{Using "add eq2"}}{H \vdash \blacksquare}}{H \vdash X \oplus Z \approx X \oplus T}$$

Sub lemma 40 $H \vdash X \oplus T \approx Y \oplus T$

Proof of sub lemma 40

$$\frac{H \vdash X \approx Y \quad \frac{\text{Using "add eq"}}{H \vdash \blacksquare}}{H \vdash X \oplus T \approx Y \oplus T}$$

Hence,

$$\frac{\frac{\text{Using 39}}{H \vdash X \oplus Z \approx X \oplus T} \quad \frac{\text{Using 39}}{H \vdash X \oplus T \approx Y \oplus T} \quad \frac{\text{Using "eq trans"}}{H \vdash \blacksquare}}{X \approx Y \wedge Z \approx T \vdash X \oplus Z \approx Y \oplus T}$$

A.3.0.5 Substraction in equality

Proof of main lemma 5 Let $H \equiv_{def} X \oplus Y \approx Z$ and $K \equiv_{def} Z \approx X \oplus Y$

Sub lemma 41 $\vdash X \oplus (Y \ominus Y) \approx X \oplus 0$

Proof of sub lemma 41

$$\frac{\frac{\text{Using "equ ref"} \quad \text{Using "opp add"}}{\vdash X \approx X \quad \vdash Y \ominus Y \approx 0} \quad \text{Using "add equ"} \quad \vdash \blacksquare}{\vdash X \approx X \wedge Y \ominus Y \approx 0}}{\vdash X \oplus (Y \ominus Y) \approx X \oplus 0}$$

Sub lemma 42 $\vdash X \oplus (Y \ominus Y) \approx X$

Proof of sub lemma 42

$$\frac{\frac{\text{Using 41}}{X \oplus (Y \ominus Y) \approx X \oplus 0} \quad \text{Using "0 add"} \quad \text{Using "= trans"} \quad \vdash \blacksquare}{\vdash X \oplus (Y \ominus Y) \approx X}}$$

Sub lemma 43 $\vdash (X \oplus Y) \ominus Y \approx X$

Proof of sub lemma 43

$$\frac{\frac{\text{Using "add assoc"}}{\vdash X \oplus (Y \ominus Y) \approx (X \oplus Y) \ominus Y} \quad \text{Using 42} \quad \text{Using "= trans"} \quad \vdash \blacksquare}{\vdash (X \oplus Y) \ominus Y \approx X}}$$

Sub lemma 44 $H \vdash (X \oplus Y) \ominus Y \approx Z \ominus Y$

Proof of sub lemma 44

$$\frac{\frac{\text{Using "equ ref"}}{H \vdash X \oplus Y \approx Z \quad H \vdash \ominus Y \approx \ominus Y} \quad \text{Using "add equ"} \quad \vdash \blacksquare}{H \vdash (X \oplus Y) \ominus Y \approx Z \ominus Y}}$$

Hence,

$$\frac{\frac{\text{Using } 43}{H \vdash (X \oplus Y) \ominus Y \approx X} \quad \frac{\text{Using } 44}{H \vdash (X \oplus Y) \ominus Y \approx Z \ominus Y}}{H \vdash (X \oplus Y) \ominus Y \approx X \wedge (X \oplus Y) \ominus Y \approx Z \ominus Y} \quad \text{Using "equ trans"} \quad \frac{}{H \vdash \blacksquare}}{X \oplus Y \approx Z \vdash X \approx Z \ominus Y}$$

and

$$\frac{\frac{K \vdash Z \approx X \oplus Y \quad \frac{\text{Using "equ ref"}}{K \vdash \blacksquare}}{K \vdash X \oplus Y \approx Z} \quad \frac{\text{Using "sub equ1"}}{K \vdash \blacksquare}}{\frac{K \vdash Z \approx Z \ominus Y}{Z \approx X \oplus Y \vdash Z \ominus Y \approx Z}} \quad \frac{\text{Using "equ ref"}}{K \vdash \blacksquare}}$$

A.3.0.6 Substraction in inequality

Proof of main lemma 6 Let $H \equiv_{def} X \oplus Y \preceq Z$ in

Sub lemma 45 $\vdash X \oplus 0 \preceq X \oplus Y \ominus Y$

Proof of sub lemma 45

$$\frac{\frac{\text{Using "opp add"} \quad \frac{\text{Using "leq equiv } \approx"}{\vdash 0 \approx Y \ominus Y} \quad \vdash \blacksquare}{\vdash 0 \preceq Y \ominus Y \wedge Y \ominus Y \preceq 0}}{\frac{\text{Using "}\preceq\text{ ref"}}{\vdash X \preceq X} \quad \frac{\vdash 0 \preceq Y \ominus Y}{\vdash X \preceq X \wedge 0 \preceq Y \ominus Y}} \quad \frac{\text{Using "}\preceq\text{ add"}}{\vdash \blacksquare}}{\vdash X \oplus 0 \preceq X \oplus Y \ominus Y}$$

Sub lemma 46 $H \vdash X \oplus Y \ominus Y \preceq Z \ominus Y$

Proof of sub lemma 46

$$\frac{\frac{H \vdash X \oplus Y \preceq Z \quad \frac{\text{"Using } \preceq\text{ ref'}}{\ominus Y \preceq \ominus Y}}{H \vdash X \oplus Y \preceq Z \wedge \ominus Y \preceq \ominus Y} \quad \frac{\text{Using "}\preceq\text{ add"}}{H \vdash \blacksquare}}{H \vdash X \oplus Y \ominus Y \preceq Z \ominus Y}$$

Sub lemma 47 $H \vdash X \oplus 0 \preceq Z \ominus Y$

Proof of sub lemma 47

$$\frac{\frac{\text{Using 46}}{H \vdash X \oplus Y \ominus Y \preceq Z \ominus Y} \quad \frac{\text{Using 45}}{H \vdash X \oplus 0 \preceq X \oplus Y \ominus Y} \quad \text{Using “}\preceq\text{ trans”}}{H \vdash X \oplus Y \ominus Y \preceq Z \ominus Y \wedge X \oplus 0 \preceq X \oplus Y \ominus Y} \quad H \vdash \blacksquare}{H \vdash X \oplus 0 \preceq Z \ominus Y}$$

Hence,

$$\frac{\frac{\text{Using 47}}{H \vdash X \oplus 0 \preceq Z \ominus Y} \quad \text{Using “}\preceq\text{ trans”}}{H \vdash X \oplus 0 \preceq Z \ominus Y \wedge X \oplus 0 \preceq X} \quad H \vdash \blacksquare}{X \oplus Y \preceq Z \vdash X \preceq Z \ominus Y}$$

A.3.0.7 Propagation of multiplication in sum

Proof of main lemma 7 By induction on $|I|$.

Let \mathcal{H}_k be the hypothesis “If $|I| \preceq k$ then $\vdash p \otimes \Sigma_{i \in I} X_i \approx \Sigma_{i \in I} p \otimes X_i$ ”

\mathcal{H}_2 ? **Sub lemma 48** $\vdash (X \oplus Y) \otimes p \approx X \otimes p \oplus Y \otimes p$

Proof of sub lemma 48

$$\frac{\frac{\text{Using “mult comm”}}{\vdash X \otimes p \approx p \otimes X} \quad \frac{\text{Using “mult comm”}}{\vdash Y \otimes p \approx p \otimes Y} \quad \text{Using “equ add”}}{\vdash X \otimes p \approx p \otimes X \wedge Y \otimes p \approx p \otimes Y} \quad \vdash \blacksquare}{\vdash X \otimes p \oplus Y \otimes p \approx p \otimes X \oplus p \otimes Y}$$

Sub lemma 49 $\vdash (X \oplus Y) \otimes p \approx p \otimes X \oplus p \otimes Y$

Proof of sub lemma 49

$$\frac{\frac{\text{Using 48}}{\vdash (X \oplus Y) \otimes p \approx X \otimes p \oplus Y \otimes p} \quad \frac{\text{Using “eq mult”}}{\vdash (X \oplus Y) \otimes p \approx X \otimes p \oplus Y \otimes p} \quad \text{Using “equ trans”}}{\vdash (X \oplus Y) \otimes p \approx X \otimes p \oplus Y \otimes p \wedge X \otimes p \oplus Y \otimes p \approx p \otimes X \oplus p \otimes Y} \quad \vdash \blacksquare}{\vdash (X \oplus Y) \otimes p \approx p \otimes X \oplus p \otimes Y}$$

Sub lemma 50 $p \otimes (X \oplus Y) \approx p \otimes X \oplus p \otimes Y$

Proof of sub lemma 50

$$\frac{\frac{\text{Using “mult comm”}}{\vdash (X \oplus Y) \otimes p \approx p \otimes X \oplus p \otimes Y} \quad \frac{\text{Using “mult comm”}}{\vdash p \otimes (X \oplus Y) \approx (X \oplus Y) \otimes p} \quad \text{Using “equ trans”}}{\vdash (X \oplus Y) \otimes p \approx p \otimes X \oplus p \otimes Y \wedge p \otimes (X \oplus Y) \approx (X \oplus Y) \otimes p} \quad \vdash \blacksquare}{\vdash p \otimes (X \oplus Y) \approx p \otimes X \oplus p \otimes Y}$$

Which proves \mathcal{H}_2 .

$\mathcal{H}_k \Rightarrow \mathcal{H}_{k \oplus 1}$? Let us suppose \mathcal{H}_k for a given k , $k \geq 2$. Let J be a set of $k+1$ integers. By definition of Σ , there is some j in J such as

$$\Sigma_{i \in J} X_i \equiv_{def} \Sigma_{i \in J \setminus \{j\}} X_i \oplus X_j$$

Hence,

$$p \otimes \Sigma_{i \in J} X_i \equiv_{def} p \otimes (\Sigma_{i \in J \setminus \{j\}} X_i \oplus X_j)$$

and

$$\Sigma_{i \in J} p \otimes X_i \equiv_{def} \Sigma_{i \in J \setminus \{j\}} p \otimes X_i \oplus p \otimes X_j$$

Let $Z \equiv_{def} \Sigma_{i \in J \setminus \{j\}} X_i$ and $T \equiv_{def} \Sigma_{i \in J \setminus \{j\}} p \otimes X_i$

Sub lemma 51 $\vdash p \otimes Z \oplus p \otimes X_j \approx T \oplus p \otimes X_j$

Proof of sub lemma 51

$$\frac{\frac{\mathcal{H}_k}{\vdash p \otimes Z \approx T} \quad \frac{\text{Using "equ ref"}}{\vdash p \otimes X_j \approx p \otimes X_j}}{\vdash p \otimes Z \approx T \wedge p \otimes X_j \approx p \otimes X_j}}{\vdash p \otimes Z \oplus p \otimes X_j \approx T \oplus p \otimes X_j}$$

Sub lemma 52 $\vdash p \otimes (\Sigma_{i \in J \setminus \{j\}} X_i \oplus X_j) \approx \Sigma_{i \in J \setminus \{j\}} p \otimes X_i \oplus p \otimes X_j$

Proof of sub lemma 52

$$\frac{\frac{\mathcal{H}_2}{\vdash p \otimes (Z \oplus X_j) \approx p \otimes Z \oplus p \otimes X_j} \quad \frac{\text{Using 51}}{\vdash p \otimes Z \oplus p \otimes X_j \approx T \oplus p \otimes X_j}}{\vdash p \otimes (Z \oplus X_j) \approx p \otimes Z \oplus p \otimes X_j \wedge p \otimes Z \oplus p \otimes X_j \approx T \oplus p \otimes X_j} \quad \frac{\text{Using "equ trans"}}{\vdash \blacksquare}}{\vdash p \otimes (\Sigma_{i \in J \setminus \{j\}} X_i \oplus X_j) \approx \Sigma_{i \in J \setminus \{j\}} p \otimes X_i \oplus p \otimes X_j}$$

which proves the hypothesis.

A.3.0.8 Variable isolation in a sum

Proof of main lemma 8 (Note : This theorem/lemma is not "static". cf. 6.2 page 27) By induction on $|I|$.

Let \mathcal{H}_k be the hypothesis "If $|I| \leq k$ then $\vdash \Sigma_{i \in I} X_i \approx \Sigma_{i \in I \setminus \{j\}} X_i \oplus X_j$ "

\mathcal{H}_2 ? $\vdash X \oplus Y \approx X \oplus Y$ and $\vdash X \oplus Y \approx Y \oplus X$

$\mathcal{H}_k \Rightarrow \mathcal{H}_{k \oplus 1}$? Let I be a set of integers of size $k+1$. By definition of Σ , there is an integer l in I such as

$$\Sigma_{i \in I} X_i \equiv_{\text{def}} \Sigma_{i \in I \setminus \{l\}} X_i \oplus X_l$$

Let $Z(p) \equiv_{\text{def}} \Sigma_{i \in I \setminus \{p\}} X_i$ and $T \equiv_{\text{def}} \Sigma_{i \in I \setminus \{j, l\}} X_i$

Sub lemma 53 $j = l \vdash Z(l) \oplus X_l \approx Z(j) \oplus X_j$

Proof of sub lemma 53

$$\frac{j = l \vdash Z(l) \approx Z(j) \quad j = l \vdash X_l \approx X_j \quad \text{Using "eq add"}}{j = l \vdash Z(l) \approx Z(j) \wedge X_l \approx X_j} \quad \frac{j = l \vdash \blacksquare}{j = l \vdash Z(l) \oplus X_l \approx Z(j) \oplus X_j}$$

Sub lemma 54 $j = l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j$

Proof of sub lemma 54

$$\frac{j = l \vdash \Sigma_{i \in I} X_i \approx Z(l) \oplus X_l \quad \frac{\text{Using 53}}{j = l \vdash Z(l) \oplus X_l \approx Z(j) \oplus X_j} \quad \text{Using "eq trans"}}{j = l \vdash \Sigma_{i \in I} X_i \approx Z(l) \oplus X_l \wedge Z(l) \oplus X_l \approx Z(j) \oplus X_j} \quad \frac{j = l \vdash \blacksquare}{j = l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j}$$

Sub lemma 55 $j \neq l \vdash \Sigma_{i \in I} X_i \approx T \oplus X_j \oplus X_l$

Proof of sub lemma 55

$$\frac{\frac{\text{Using } \mathcal{H}_k}{j \neq l \vdash Z(l) \approx T \oplus X_j} \quad \frac{\text{Using "add eq"}}{j \neq l \vdash \blacksquare}}{j \neq l \vdash Z(l) \oplus X_l \approx T \oplus X_j \oplus X_l} \quad \frac{j \neq l \vdash \Sigma_{i \in I} X_i \approx Z(l) \oplus X_l \quad \text{Using "eq trans"}}{j \neq l \vdash \Sigma_{i \in I} X_i \approx Z(l) \oplus X_l \wedge Z(l) \oplus X_l \approx T \oplus X_j \oplus X_l} \quad \frac{j \neq l \vdash \blacksquare}{j \neq l \vdash \Sigma_{i \in I} X_i \approx T \oplus X_j \oplus X_l}$$

Sub lemma 56 $\vdash X \oplus (Y \oplus T) \approx X \oplus (T \oplus Y)$

Proof of sub lemma 56

$$\frac{\frac{\text{Using "eq ref"}}{\vdash X \approx X} \quad \frac{\text{Using "add comm"}}{Y \oplus T \approx T \oplus Y}}{\vdash X \approx X \wedge Y \oplus T \approx T \oplus Y} \quad \frac{\text{Using "equ add"}}{\vdash \blacksquare}}{\vdash X \oplus (Y \oplus T) \approx X \oplus (T \oplus Y)}$$

Sub lemma 57 $\vdash X \oplus (Y \oplus T) \approx (X \oplus T) \oplus Y$

Proof of sub lemma 57

$$\frac{\frac{\text{Using 56}}{\vdash X \oplus (T \oplus Y) \approx X \oplus (Y \oplus T)} \quad \frac{\text{Using "add assoc"}}{\vdash (X \oplus T) \oplus Y \approx X \oplus (Y \oplus T)}}{\vdash X \oplus (T \oplus Y) \approx X \oplus (Y \oplus T) \wedge (X \oplus T) \oplus Y \approx X \oplus (Y \oplus T)} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{\vdash X \oplus (Y \oplus T) \approx (X \oplus T) \oplus Y}$$

Sub lemma 58 $\vdash X \oplus Y \oplus T \approx X \oplus T \oplus Y$

Proof of sub lemma 58

$$\frac{\frac{\text{Using "add assoc"}}{\vdash (X \oplus Y) \oplus T \approx X \oplus (Y \oplus T)} \quad \frac{\text{Using 57}}{\vdash X \oplus (Y \oplus T) \approx (X \oplus T) \oplus Y}}{\vdash (X \oplus Y) \oplus T \approx X \oplus (Y \oplus T) \wedge X \oplus (Y \oplus T) \approx (X \oplus T) \oplus Y} \quad \frac{\text{Using "eq trans"}}{\vdash \blacksquare}}{\vdash (X \oplus Y) \oplus T \approx (X \oplus T) \oplus Y}$$

Sub lemma 59 $j \neq l \vdash T \oplus X_j \oplus X_l \approx Z(j) \oplus X_j$

Proof of sub lemma 59

$$\frac{\frac{\text{Using 58}}{\vdash (T \oplus X_j) \oplus X_l \approx (T \oplus X_l) \oplus X_j} \quad \vdash (T \oplus X_l) \oplus X_j \approx Z(j) \oplus X_j}{j \neq l \vdash (T \oplus X_j) \oplus X_l \approx (T \oplus X_l) \oplus X_j \wedge (T \oplus X_l) \oplus X_j \approx Z(j) \oplus X_j} \quad \frac{\text{Using "eq trans"}}{j \neq l \vdash \blacksquare}}{j \neq l \vdash T \oplus X_j \oplus X_l \approx Z(j) \oplus X_j}$$

Sub lemma 60 $j \neq l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j$

Proof of sub lemma 60

$$\frac{\frac{\text{Using 55}}{j \neq l \vdash \Sigma_{i \in I} X_i \approx T \oplus X_j \oplus X_l} \quad \frac{\text{Using 59}}{T \oplus X_j \oplus X_l \approx Z(j) \oplus X_j}}{j \neq l \vdash \Sigma_{i \in I} X_i \approx T \oplus X_j \oplus X_l \wedge T \oplus X_j \oplus X_l \approx Z(j) \oplus X_j} \quad \frac{\text{Using "eq trans"}}{j \neq l \vdash \blacksquare}}{j \neq l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j}$$

Hence,

$$\frac{\frac{\text{Using 60}}{j \neq l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j} \quad \frac{\text{Using 54}}{j = l \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j}}{(j = l \vee j \neq l) \vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j}}{\vdash \Sigma_{i \in I} X_i \approx Z(j) \oplus X_j}$$

Which proves the hypothesis.

A.3.0.9 Variable resolution

Proof of main lemma 9 Let $A \equiv_{def} \sum_{i \in I \setminus \{k\}} X_i$, $B \equiv_{def} \sum_{i \in I \setminus \{k\}} \ominus X_i$ and $H \equiv_{def} \sum_{i \in I} X_i \approx c$ in

Sub lemma 61 $H \vdash A \oplus X_k \approx c$

Proof of sub lemma 61

$$\frac{\frac{\text{Using "var isol"}}{H \vdash A \oplus X_k \approx \sum_{i \in I} X_i} \quad H \vdash \sum_{i \in I} X_i \approx c \text{ Using "equ trans"}}{H \vdash A \oplus X_k \approx \sum_{i \in I} X_i \wedge \sum_{i \in I} X_i \approx c} \quad H \vdash \blacksquare}{H \vdash A \oplus X_k \approx c}$$

Sub lemma 62 $H \vdash X_k \approx c \ominus A$

Proof of sub lemma 62

$$\frac{\frac{\text{Using "add comm"}}{H \vdash X_k \oplus A \approx A \oplus X_k} \quad \frac{\text{Using 61}}{H \vdash A \oplus X_k \approx c} \quad \text{Using "add trans"}}{H \vdash X_k \oplus A \approx A \oplus X_k \wedge A \oplus X_k \approx c} \quad H \vdash \blacksquare}{\frac{H \vdash X_k \oplus A \approx c}{H \vdash X_k \approx c \ominus A}} \quad \text{Using "sub eq"} \quad H \vdash \blacksquare$$

Hence,

$$\frac{\frac{\text{Using 62}}{H \vdash X_k \approx c \ominus A} \quad \frac{\frac{\text{Using "mult sum"}}{H \vdash c \approx c} \quad \frac{H \vdash \ominus A \approx B}{H \vdash c \approx c \wedge \ominus A \approx B}}{\text{Using "equ add"} \quad H \vdash \blacksquare}}{H \vdash X_k \approx c \ominus A \wedge c \ominus A \approx c \oplus B} \quad \text{Using "equ add"} \quad H \vdash \blacksquare}{\sum_{i \in I} X_i \approx c \vdash X_k \approx c \oplus \sum_{i \in I \setminus \{k\}} \ominus X_i}$$

A.3.0.10 0 Addition

Proof of main lemma 10

$$\frac{\text{Using "0 add"} \quad \text{Using "eq ref"}}{\frac{\vdash X \oplus 0 \approx X}{\vdash X \approx X \oplus 0}} \quad \vdash \blacksquare$$

A.4 The Omega test

A.4.0.11 Equality normalization

Proof of main lemma 11 (Note : This theorem/lemma is not “static”. cf. 6.2 page 27) Let $S \equiv_{def} \sum_{i \in [1, n]} a_i \otimes X_i$, $T \equiv_{def} \sum_{i \in [1, n]} \frac{a_i}{p} \otimes X_i$ and $U \equiv_{def} \sum_{i \in [1, n]} p \otimes \frac{a_i}{p} \otimes X_i$. Let us suppose $p|a_1, p|a_2, \dots, p|a_n$.

Sub lemma 63 $\vdash S \approx U$

Proof of sub lemma 63 DDT: Trivial but long. To do.

Sub lemma 64 $\vdash S \approx p \otimes T$

Proof of sub lemma 64

$$\frac{\frac{\text{Using 63}}{\vdash S \approx U} \quad \frac{\text{Using “mult sum”}}{\vdash U \approx p \otimes T}}{\vdash S \approx U \wedge U \approx p \otimes T}}{\vdash S \approx p \otimes T}$$

Hence,

$$\frac{\frac{\frac{\text{Using 64}}{\vdash S \approx p \otimes T} \quad \frac{\frac{\text{Using “0 add”}}{\vdash p \otimes T \oplus 0 \approx p \otimes T} \quad \frac{\text{Using “eq ref”}}{\vdash \blacksquare}}{\vdash p \otimes T \approx p \otimes T \oplus 0}}{\vdash S \approx p \otimes T \wedge p \otimes T \approx p \otimes T \oplus 0}}{\vdash S \approx p \otimes T \oplus 0} \quad \vdash 0 \leq 0}{\frac{\frac{\vdash 0 \leq 0 \wedge 0 < p \wedge S \approx p \otimes T \oplus 0}{\vdash \exists b, 0 \leq b \wedge b < p \wedge S \approx p \otimes T \oplus b}}{p|a_1 \wedge p|a_2 \wedge \dots \wedge p|a_n \vdash (\sum_{i \in [1, n]} a_i \otimes X_i) \setminus p \approx \sum_{i \in [1, n]} \frac{a_i}{p} \otimes X_i}}$$

Proof of algorithm step 1 (Note : This theorem/lemma is not “static”. cf. 6.2 page 27) Let $H \equiv_{def} \sum_{i \in [1, n]} a_i \otimes X_i \approx c$, $S \equiv_{def} \sum_{i \in [1, n]} a_i \otimes X_i$ and $T \equiv_{def} \sum_{i \in [1, n]} \frac{a_i}{p} \otimes X_i$. Let us suppose $p|a_1, p|a_2, \dots, p|a_n$.

Sub lemma 65 $H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor$

Proof of sub lemma 65

$$\begin{array}{c}
\text{Using "mult on } \mathbb{Z}\text{"} \\
\frac{H \vdash c \approx p \cdot \lfloor \frac{c}{p} \rfloor}{H \vdash c \approx p \cdot \lfloor \frac{c}{p} \rfloor \wedge p \cdot \lfloor \frac{c}{p} \rfloor \approx p \otimes \lfloor \frac{c}{p} \rfloor} \\
\frac{H \vdash S \approx c}{H \vdash S \approx c \wedge c \approx p \otimes \lfloor \frac{c}{p} \rfloor} \\
\hline
\frac{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor}{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor} \quad \text{Using "eq trans"}
\end{array}$$

Sub lemma 66 $H \vdash S \setminus p \approx \lfloor \frac{c}{p} \rfloor$

Proof of sub lemma 66

$$\begin{array}{c}
\text{Using 65} \quad \text{Using "0 add2"} \\
\frac{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor}{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor \wedge p \otimes \lfloor \frac{c}{p} \rfloor \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0} \\
\frac{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0}{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0} \\
\frac{H \vdash 0 \quad H \vdash b < p}{H \vdash 0 \leq 0 \wedge b < p \wedge S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0} \\
\frac{H \vdash \exists b, 0 \leq b \wedge b < p \wedge S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus b}{H \vdash S \setminus p \approx \lfloor \frac{c}{p} \rfloor}
\end{array}$$

$$\begin{array}{c}
\text{Using "div sum"} \quad \text{Using "eq ref"} \\
\frac{H \vdash S \setminus p \approx T}{H \vdash T \approx S \setminus p} \quad \frac{H \vdash \blacksquare}{H \vdash S \setminus p \approx \lfloor \frac{c}{p} \rfloor} \quad \text{Using 66} \\
\hline
\frac{H \vdash T \approx S \setminus p \wedge S \setminus p \approx \lfloor \frac{c}{p} \rfloor}{H \vdash T \approx \lfloor \frac{c}{p} \rfloor}
\end{array}$$

A.4.0.12 Unsatisfiable equality

Proof of algorithm step 2 (Note : This theorem/lemma is not "static". cf. 6.2 page 27) Let $H \equiv_{def} p|a_1 \wedge p|a_2 \wedge \dots \wedge p|a_n \wedge \sum_{i=1}^{i=n} a_i \otimes X_i \approx c$ and $S \equiv_{def} \sum_{i=1}^{i=n} a_i \otimes X_i$.

Sub lemma 67 $H \vdash S \% p \approx 0$

Proof of sub lemma 67

$$\begin{array}{c}
\frac{\text{Using } 64}{H \vdash S \approx p \otimes S \setminus p} \quad \frac{\text{Using "0 add2"}}{H \vdash p \otimes S \setminus p \approx p \otimes S \setminus p \oplus 0} \\
\frac{H \vdash S \approx p \otimes S \setminus p \wedge p \otimes S \setminus p \approx p \otimes S \setminus p \oplus 0}{H \vdash S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0} \\
\frac{\vdash 0 \leq 0 \quad \vdash 0 < p}{H \vdash 0 \leq 0 \wedge 0 < p \wedge S \approx p \otimes \lfloor \frac{c}{p} \rfloor \oplus 0} \\
\frac{H \vdash \exists Y, 0 \leq 0 \wedge 0 < p \wedge S \approx p \otimes Y \oplus 0}{H \vdash S \% p \approx 0}
\end{array}$$

Sub lemma 68 $H \vdash c \approx p \otimes S \setminus p$

Proof of sub lemma 68

$$\begin{array}{c}
\frac{\text{Using } 64}{H \vdash S \approx c \quad H \vdash p \otimes S \setminus p \approx S} \quad \frac{\text{Using "eq trans"}}{H \vdash p \otimes S \setminus p \approx S \wedge S \approx c} \quad \frac{H \vdash \blacksquare}{\text{Using "eq ref"}} \\
\frac{H \vdash p \otimes S \setminus p \approx c}{H \vdash c \approx p \otimes S \setminus p}
\end{array}$$

Sub lemma 69 $H \vdash c \% p \approx 0$

Proof of sub lemma 69

$$\begin{array}{c}
\frac{\text{Using } 68}{H \vdash c \approx p \otimes S \setminus p} \quad \frac{\text{Using "0 add2"}}{p \otimes S \setminus p \approx p \otimes S \setminus p \oplus 0} \quad \frac{\text{Using "eq trans"}}{H \vdash c \approx p \otimes S \setminus p \wedge p \otimes S \setminus p \approx p \otimes S \setminus p \oplus 0} \\
\frac{H \vdash 0 \leq 0 \quad H \vdash 0 < p}{H \vdash c \approx S \setminus p \oplus 0} \\
\frac{H \vdash 0 \leq 0 \wedge 0 < p \wedge c \approx p \otimes S \setminus p \oplus 0}{H \vdash \exists Y, 0 \leq 0 \wedge 0 < p \wedge c \approx p \otimes Y \oplus 0} \\
\frac{H \vdash \exists Y, 0 \leq 0 \wedge 0 < p \wedge c \approx p \otimes Y \oplus 0}{H \vdash c \% p \approx 0}
\end{array}$$

Hence,

$$\begin{array}{c}
\frac{\text{Using } 69}{H \vdash c \% p \approx 0} \quad \frac{\text{Using "isdiv in } \mathbb{Z}}{H \vdash p \parallel c} \quad \frac{H \vdash \blacksquare}{H \vdash p | c}
\end{array}$$

A.4.0.13 Inequality tightening

Proof of algorithm step 3 (Note : This theorem/lemma is not “static”. cf. 6.2 page 27) Let $H \equiv_{def} 0 \leq k \wedge c \approx \sum_{i \in I} a_i \otimes X_i \oplus k$, $S \equiv_{def} \sum_{i \in I} a_i \otimes X_i$, $T \equiv_{def} \sum_{i \in I} \frac{a_i}{p} \otimes X_i$ and $d \equiv_{def} \lfloor \frac{c}{p} \rfloor - \lfloor \frac{c-k}{p} \rfloor$. Let us suppose $p|a_1, p|a_2, \dots, p|a_n$.

Sub lemma 70 $H \vdash S \approx p \otimes (\lfloor \frac{c}{p} \rfloor \ominus r) \oplus d$

Proof of sub lemma 70

$$\frac{\frac{\frac{H \vdash c \approx S \oplus k \quad \text{Using “eq sub”}}{H \vdash c \ominus k \approx S} \quad H \vdash \blacksquare}{H \vdash S \approx c \ominus k} \quad \frac{\text{Using “eq ref”}}{H \vdash \blacksquare}}{H \vdash \blacksquare} \quad \text{Using “eq norm”}}{H \vdash T \approx \lfloor \frac{c \ominus k}{p} \rfloor} \\ \frac{H \vdash T \approx \lfloor \frac{c}{p} \rfloor \ominus d}{H \vdash T \oplus d \approx \lfloor \frac{c}{p} \rfloor}$$

Hence,

$$\frac{\frac{\text{Using 70}}{H \vdash T \oplus d \approx \lfloor \frac{c}{p} \rfloor} \quad \frac{\text{Using “eq ref”}}{H \vdash \blacksquare}}{H \vdash \lfloor \frac{c}{p} \rfloor \approx T \oplus d} \quad H \vdash 0 \leq d}{\frac{H \vdash 0 \leq d \wedge \lfloor \frac{c}{p} \rfloor \approx T \oplus d}{H \vdash \exists a, 0 \leq a \wedge \lfloor \frac{c}{p} \rfloor \approx T \oplus a}} \\ H \vdash T \preceq \lfloor \frac{c}{p} \rfloor$$

A.4.0.14 Real shadow

Proof of main lemma 12 Let $H \equiv_{def} 0 \leq t \wedge Y \approx X \oplus t \wedge 0 \leq Z \wedge Z \approx z \oplus 0$

Sub lemma 71 $H \vdash Z \otimes t \approx z \cdot t$

Proof of sub lemma 71

$$\frac{\frac{\text{Using “mult eq”}}{H \vdash Z \approx z} \quad H \vdash \blacksquare}{H \vdash Z \otimes t \approx z \otimes t} \quad \frac{\text{Using “mult on } \mathbb{Z} \text{”}}{H \vdash z \otimes t \approx z \cdot t}}{H \vdash Z \otimes t \approx z \otimes t \wedge z \otimes t \approx z \cdot t} \\ H \vdash Z \otimes t \approx z \cdot t$$

Sub lemma 72 $H \vdash Z \otimes (X \oplus t) \approx Z \otimes X \oplus z \cdot t$

Proof of sub lemma 72

$$\frac{\frac{\text{Using "mult/add dist"}}{H \vdash Z \otimes (X \oplus t) \approx (Z \otimes X) \oplus (Z \otimes t)} \quad \frac{\frac{\text{Using 71}}{H \vdash Z \otimes t \approx z \cdot t} \quad \frac{\text{Using "add eq"}}{\vdash \blacksquare}}{H \vdash (Z \otimes X) \oplus (Z \otimes t) \approx Z \otimes X \oplus z \cdot t}}{H \vdash Z \otimes (X \oplus t) \approx (Z \otimes X) \oplus (Z \otimes t) \wedge (Z \otimes X) \oplus (Z \otimes t) \approx Z \otimes X \oplus z \cdot t}}{H \vdash Z \otimes (X \oplus t) \approx Z \otimes X \oplus z \cdot t}$$

Sub lemma 73 $H \vdash Z \otimes Y \approx Z \otimes X \oplus z \cdot t$

Proof of sub lemma 73

$$\frac{\frac{\frac{\text{Using "mult eq"}}{H \vdash Y \approx X \oplus t} \quad \frac{H \vdash \blacksquare}}{H \vdash Z \otimes Y \approx Z \otimes (X \oplus t)} \quad \frac{\text{Using 72}}{H \vdash Z \otimes (X \oplus t) \approx Z \otimes X \oplus z \cdot t}}{H \vdash Z \otimes Y \approx Z \otimes (X \oplus t) \wedge Z \otimes (X \oplus t) \approx Z \otimes X \oplus z \cdot t}}{H \vdash Z \otimes Y \approx Z \otimes X \oplus z \cdot t}$$

Hence,

$$\frac{\frac{\text{Using 71}}{H \vdash Z \otimes Y \approx Z \otimes X \oplus z \cdot t} \quad H \vdash 0 \leq z}{H \vdash 0 \leq z \cdot t \wedge Z \otimes Y \approx Z \otimes X \oplus z \cdot t}}{\exists z, t, H \vdash \exists a, 0 \leq a \wedge Z \otimes Y \approx Z \otimes X \oplus a}{X \preceq Y \wedge 0 \preceq Z \vdash Z \otimes X \preceq Z \otimes Y}$$

Proof of algorithm step 4 Let $H \equiv_{def} a \otimes Z \preceq A \wedge B \preceq b \otimes Z$

Sub lemma 74 $H \vdash a \otimes B \preceq a \otimes (b \otimes Z)$

Proof of sub lemma 74

$$\frac{\frac{H \vdash 0 \leq a}{H \vdash 0 \preceq a} \quad H \vdash B \preceq b \otimes Z \quad \text{Using "mult ineq"}}{H \vdash 0 \preceq a \wedge B \preceq b \otimes Z} \quad \frac{H \vdash \blacksquare}{H \vdash a \otimes B \preceq a \otimes (b \otimes Z)}$$

Sub lemma 75 $H \vdash (a \otimes b) \otimes Z \preceq a \cdot b \otimes Z$

Proof of sub lemma 75

$$\frac{\frac{\text{Using "mult on } \mathbb{Z}\text{"}}{H \vdash (a \otimes b) \approx b \otimes Z}}{H \vdash (a \otimes b) \otimes Z \approx a \cdot b \otimes Z} \quad \frac{\text{Using "lt as leq"} \quad H \vdash \blacksquare}{H \vdash (a \otimes b) \otimes Z \preceq a \cdot b \otimes Z}}$$

Sub lemma 76 $H \vdash a \otimes (b \otimes Z) \preceq a \cdot b \otimes Z$

Proof of sub lemma 76

$$\frac{\frac{\text{Using 75}}{H \vdash (a \otimes b) \otimes Z \preceq a \cdot b \otimes Z} \quad \frac{\frac{\text{Using "mult assoc"} \quad \text{Using "lt as leq"}}{H \vdash a \otimes (b \otimes Z) \approx (a \otimes b) \otimes Z} \quad H \vdash \blacksquare}{H \vdash a \otimes (b \otimes Z) \preceq (a \otimes b) \otimes Z}}{H \vdash (a \otimes b) \otimes Z \preceq a \cdot b \otimes Z \wedge a \otimes (b \otimes Z) \preceq (a \otimes b) \otimes Z} \quad \frac{\text{Using "leq trans"} \quad H \vdash \blacksquare}{H \vdash a \otimes (b \otimes Z) \preceq a \cdot b \otimes Z}}$$

Sub lemma 77 $H \vdash a \otimes B \preceq a \cdot b \otimes Z$

Proof of sub lemma 77

$$\frac{\frac{\text{Using 74}}{H \vdash a \otimes B \preceq a \otimes (b \otimes Z)} \quad \frac{\text{Using 76}}{H \vdash a \otimes (b \otimes Z) \preceq a \cdot b \otimes Z}}{H \vdash a \otimes B \preceq a \otimes (b \otimes Z) \wedge a \otimes (b \otimes Z) \preceq a \cdot b \otimes Z} \quad \frac{\text{Using "leq trans"} \quad H \vdash \blacksquare}{H \vdash a \otimes B \preceq a \cdot b \otimes Z}}$$

Sub lemma 78 $H \vdash a \cdot b \otimes Z \preceq b \otimes A$

Proof of sub lemma 78 *The proof is very similar to that of lemma 77*

$$\frac{\frac{\text{Using 77}}{H \vdash a \otimes B \preceq a \cdot b \otimes Z} \quad \frac{\text{Using 78}}{H \vdash a \cdot b \otimes Z \preceq b \otimes A}}{H \vdash a \otimes B \preceq a \cdot b \otimes Z \wedge a \cdot b \otimes Z \preceq b \otimes A} \quad \frac{}{a \otimes Z \preceq A \wedge B \preceq b \otimes Z \vdash a \otimes B \preceq b \otimes A}}$$

A.4.0.15 Exhaustive check

Proof of algorithm step 5 (Note : This theorem/lemma is not "static". cf. 6.2 page 27) Let $H \equiv_{def} 0 \leq l \wedge (a-1) \cdot (b-1) \approx b \otimes A \ominus a \otimes B \oplus l \wedge 0 \leq k \wedge A \approx a \otimes \xi \oplus k \wedge 0 \leq m \wedge b \otimes \xi \approx B \oplus m$. Suppose $1 \leq a \wedge 1 \leq b$.

Sub lemma 79 $H \vdash a \otimes m \oplus a \otimes B \approx a \cdot m \oplus a \otimes B$

Proof of sub lemma 79

$$\frac{\frac{\text{Using "mult on } \mathbb{Z}"}{\vdash a \otimes m \approx a \cdot m} \quad \frac{\text{Using "eq mult"}}{\vdash \blacksquare}}{\vdash a \otimes m \oplus a \otimes B \approx a \cdot m \oplus a \otimes B}}$$

Sub lemma 80 $H \vdash B \oplus m \preceq b \otimes \xi$

Proof of sub lemma 80

$$\frac{H \vdash B \oplus m \approx b \otimes \xi \quad \frac{\text{Using "eq as ineq"}}{H \vdash \blacksquare}}{H \vdash B \oplus m \preceq b \otimes \xi}}$$

Sub lemma 81 $H \vdash m \oplus B \preceq b \otimes \xi$

Proof of sub lemma 81

$$\frac{\frac{\frac{\text{Using "add comm"} \quad \frac{\text{Using "eq as ineq"}}{\vdash m \oplus B \approx B \oplus m} \quad \vdash \blacksquare}}{H \vdash m \oplus B \preceq B \oplus m} \quad \frac{\text{Using 80}}{H \vdash B \oplus m \preceq b \otimes \xi} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash m \oplus B \preceq B \oplus m \wedge B \oplus m \preceq b \otimes \xi} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash m \oplus B \preceq b \otimes \xi}}$$

Sub lemma 82 $H \vdash a \otimes (m \oplus B) \preceq a \otimes b \otimes \xi$

Proof of sub lemma 82

$$\frac{\frac{\frac{\text{Using 81}}{H \vdash m \oplus B \preceq b \otimes \xi} \quad \frac{\frac{\text{Using "leq on } \mathbb{Z}"}{H \vdash 0 \preceq a} \quad H \vdash \blacksquare}}{H \vdash 0 \preceq a} \quad \frac{\text{Using "eq mult"}}{H \vdash \blacksquare}}{H \vdash m \oplus B \preceq b \otimes \xi \wedge 0 \preceq a} \quad \frac{\text{Using "eq mult"}}{H \vdash \blacksquare}}{H \vdash a \otimes (m \oplus B) \preceq a \otimes b \otimes \xi}}$$

Sub lemma 83 $H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes (m \oplus B)$

Proof of sub lemma 83

$$\frac{\frac{\text{Using "add/mult dist"}}{H \vdash a \otimes m \oplus a \otimes B \approx a \otimes (m \oplus B)} \quad \frac{\text{Using "eq as ineq"}}{H \vdash \blacksquare}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes (m \oplus B)}}$$

Sub lemma 84 $H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes b \otimes \xi$

Proof of sub lemma 84

$$\frac{\frac{\text{Using 83}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes (m \oplus B)} \quad \frac{\text{Using 82}}{H \vdash a \otimes (m \oplus B) \preceq a \otimes b \otimes \xi}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes (m \oplus B) \wedge a \otimes (m \oplus B) \preceq a \otimes b \otimes \xi} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes b \otimes \xi}$$

Sub lemma 85 $H \vdash a \cdot m \oplus a \otimes B \preceq a \otimes b \otimes \xi$

Proof of sub lemma 85

$$\frac{\frac{\text{Using 79}}{\vdash a \otimes m \oplus a \otimes B \approx a \cdot m \oplus a \otimes B} \quad \frac{\text{Using "eq as ineq"}}{\vdash \blacksquare}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \cdot m \oplus a \otimes B} \quad \frac{\text{Using 84}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \otimes b \otimes \xi} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash a \otimes m \oplus a \otimes B \preceq a \cdot m \oplus a \otimes B \wedge a \cdot m \oplus a \otimes B \preceq a \otimes b \otimes \xi} \quad \frac{}{H \vdash a \cdot m \oplus a \otimes B \preceq a \otimes b \otimes \xi}$$

Sub lemma 86 $\vdash a \cdot b \otimes \xi \preceq a \otimes b \otimes \xi$

Proof of sub lemma 86

$$\frac{\frac{\text{Using "mult on } \mathbb{Z}}{\vdash a \cdot b \approx a \otimes b} \quad \frac{\text{Using "eq mult"}}{\vdash \blacksquare}}{\vdash a \cdot b \otimes \xi \approx a \otimes b \otimes \xi} \quad \frac{\text{Using "eq as ineq"}}{H \vdash \blacksquare}}{\vdash a \cdot b \otimes \xi \preceq a \otimes b \otimes \xi}$$

Sub lemma 87 $H \vdash m \cdot a \oplus a \otimes B \preceq a \cdot b \otimes \xi$

Proof of sub lemma 87

$$\frac{\frac{\text{Using 85}}{H \vdash a \cdot m \oplus a \otimes B \preceq a \otimes b \otimes \xi} \quad \frac{\text{Using 86}}{H \vdash a \cdot b \otimes \xi \preceq a \otimes b \otimes \xi}}{H \vdash a \cdot m \oplus a \otimes B \preceq a \otimes b \otimes \xi \wedge a \cdot b \otimes \xi \preceq a \otimes b \otimes \xi} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash a \cdot m \oplus a \otimes B \preceq a \cdot b \otimes \xi}$$

Sub lemma 88 $H \vdash b \otimes (a \otimes \xi) \preceq b \otimes A$

Proof of sub lemma 88

$$\frac{H \vdash a \otimes \xi \preceq A \quad \frac{H \vdash 0 \leq b \quad \frac{\text{Using "leq on } \mathbb{Z}"}{H \vdash \blacksquare}}{H \vdash 0 \preceq b}}{H \vdash a \otimes \xi \preceq A \wedge 0 \preceq b} \quad \frac{\text{Using "ineq mult"}}{H \vdash \blacksquare}}{H \vdash b \otimes (a \otimes \xi) \preceq b \otimes A}$$

Sub lemma 89 $H \vdash a \otimes b \otimes \xi \preceq b \otimes A$

Proof of sub lemma 89

$$\frac{\frac{\text{Using 88}}{H \vdash b \otimes (a \otimes \xi) \preceq b \otimes A} \quad \frac{\frac{\text{Using "mult assoc"}}{H \vdash b \otimes a \otimes \xi \approx b \otimes (a \otimes \xi)} \quad \frac{\text{Using "eq as ineq"}}{H \vdash \blacksquare}}{H \vdash b \otimes a \otimes \xi \preceq b \otimes (a \otimes \xi)}}{H \vdash b \otimes (a \otimes \xi) \preceq b \otimes A \wedge b \otimes a \otimes \xi \preceq b \otimes (a \otimes \xi)} \quad \frac{\text{Using 88}}{H \vdash b \otimes a \otimes \xi \preceq b \otimes A}}{H \vdash b \otimes a \otimes \xi \preceq b \otimes A}$$

Sub lemma 90 $H \vdash a \cdot b \otimes \xi \preceq b \otimes A$

Proof of sub lemma 90

$$\frac{\frac{\text{Using 86}}{H \vdash a \cdot b \otimes \xi \preceq a \otimes b \otimes \xi} \quad \frac{\text{Using 89}}{H \vdash b \otimes a \otimes \xi \preceq b \otimes A} \quad \frac{\text{Using "leq trans"}}{H \vdash \blacksquare}}{H \vdash b \cdot a \otimes \xi \preceq b \otimes a \otimes \xi \wedge b \otimes a \otimes \xi \preceq b \otimes A} \quad \frac{\text{Using 86}}{H \vdash a \cdot b \otimes \xi \preceq b \otimes A}}$$

Sub lemma 91 $H \vdash m \cdot a \preceq b \otimes A \ominus a \otimes B$

Proof of sub lemma 91

$$\frac{\frac{\text{Using 87}}{H \vdash m \cdot a \oplus a \otimes B \preceq a \cdot b \otimes \xi} \quad \frac{\text{Using 90}}{H \vdash a \cdot b \otimes \xi \preceq b \otimes A}}{H \vdash m \cdot a \oplus a \otimes B \preceq a \cdot b \otimes \xi \wedge a \cdot b \otimes \xi \preceq b \otimes A} \quad \frac{\text{Using "sub eq"}}{H \vdash \blacksquare}}{H \vdash m \cdot a \oplus a \otimes B \preceq b \otimes A} \quad \frac{\text{Using 90}}{H \vdash m \cdot a \preceq b \otimes A \ominus a \otimes B}$$

Sub lemma 92 $H \vdash m \cdot a \preceq (a \cdot b - a - b)$

Proof of sub lemma 92

$$\frac{\frac{\frac{H \vdash 0 \leq l \quad H \vdash (a \cdot b - a - b) \approx b \otimes A \ominus a \otimes B \oplus l}{H \vdash 0 \leq l \wedge (a \cdot b - a - b) \approx b \otimes A \ominus a \otimes B \oplus l}}{\text{Using 91}} \quad \frac{H \vdash \exists n, 0 \leq n \wedge (a \cdot b - a - b) \approx b \otimes A \ominus a \otimes B \oplus n}{H \vdash b \otimes A \ominus a \otimes B \preceq (a \cdot b - a - b)}}{H \vdash m \cdot a \preceq b \otimes A \ominus a \otimes B} \quad \frac{H \vdash m \cdot a \preceq b \otimes A \ominus a \otimes B \wedge b \otimes A \ominus a \otimes B \preceq (a \cdot b - a - b)}{H \vdash m \cdot a \preceq (a \cdot b - a - b)}$$

Hence,

$$\frac{\frac{\frac{\text{Using 92}}{H \vdash m \cdot a \preceq (a \cdot b - a - b)} \quad \frac{\text{Using "leq on } \mathbb{Z}"}{H \vdash \blacksquare}}{H \vdash 0 \leq m} \quad \frac{H \vdash m \cdot a \leq (a \cdot b - a - b)}{H \vdash 0 \leq m \wedge m \cdot a \leq (a \cdot b - a - b)} \quad H \vdash b \otimes \xi \approx B \oplus m}{H \vdash \exists i, 0 \leq i \wedge i \cdot a \leq (a \cdot b - a - b) \wedge b \otimes \xi \approx B \oplus i}}$$

B Proofs on Solver2FOL

B.1 Expression through Disjunctive Normal Form

Property 2 Let $] - R, R]$ be the range of representable integers for a given computer.

Let us consider function $t : x \mapsto 2^x$

In Solver2FOL's language, any expression of function f as a Disjunctive Normal Form (DNF) which is exact on $[0, R]$ will require at least $R/4$ elementary conjunctions.

Proof of property 2 Suppose that t can be represented on $[0, R]$ as

$$T(x, y) \equiv_{def} \bigvee_{i=1}^{i=n} C_i(x, y)$$

where

$$T(x, y) \equiv y = t(x)$$

and $C_i(x, y)$ is a conjunction of elementary (linear) constraints.

Suppose $n < R/4$.

This means that at least one $C_i(x, y)$ will hold a constraint which will be valid for at least 3 values of x . Additionally, since $C_i(x, y)$ is in normal form, this means that these constraints define an integer interval and that the value of y will be defined by the same formula.

In other words,

$$\begin{cases} y &= a \cdot x + b \\ 2 \cdot y &= a \cdot (x + 1) + b \\ 4 \cdot y &= a \cdot (x + 2) + b \end{cases}$$

A resolution of this system gives :

$$a = 0, b = 0, y = 0$$

Since all values of x considered are in $[0, R]$, we necessarily have $y > 0$. Hence, contradiction.

Since $n > R/4$, there are at least $R/4$ elementary conjunctions.